arXiv:0812.0228v1 [hep-th] 1 Dec 2008

Gauge Orbit Types for Theories with Classical Compact Gauge Group

A. Hertsch, G. Rudolph and M. Schmidt
Institut für Theoretische Physik, Universität Leipzig
Augustusplatz 10/11, 04109 Leipzig, Germany

December 1, 2008

Abstract

We determine the orbit types of the action of the group of local gauge transformations on the space of connections in a principal bundle with structure group O(n), SO(n) or Sp(n) over a closed, simply connected manifold of dimension 4. Complemented with earlier results on U(n) and SU(n) this completes the classification of the orbit types for all classical compact gauge groups over such space-time manifolds. On the way we derive the classification of principal bundles with structure group SO(n) over these manifolds and the Howe subgroups of SO(n).

Contents

1 Introduction 2

2 Orbit types and holonomy-induced Howe subbundles 4

3 Howe subgroups 6
3.1 Howe subgroups of O(n) . . . 6
3.2 Howe subgroups of Sp(n) . . . 8
3.3 Howe subgroups of SO(n) . . . 9

4 Principal bundles with a Howe subgroup as structure group 16
4.1 Principal bundles with structure group O(n) or SO(n) . . . 17
4.2 Principal bundles with structure groups H_0, H and S$H^\pm$ . . . 27

5 Howe subbundles 30
5.1 Howe subbundles of principal O(n)- and SO(n)-bundles . . . 30
5.2 Howe subbundles of principal Sp(n)-bundles . . . 33

6 Holonomy-induced Howe subbundles 34

7 Factorization by the action of the structure group 36
7.1 Structure groups O(n) and Sp(n) . . . 36
7.2 Structure group SO(n) . . . 38

8 Summary: Classification of orbit types 40

9 Examples 42
9.1 Sp(1) and Sp(2) . . . 43
9.2 O(4) . . . 45
9.3 SO(4) . . . 47

A Universal bundles 49

B Eilenberg-MacLane spaces 50

C Cohomology operations 51

D Characteristic classes 52

E The Moore-Postnikov tower 54

1 Introduction

The principle of local gauge invariance plays a fundamental role in modern theoretical physics. Its application to the theory of particle interactions gave rise to the standard model, which proved to be successful from both theoretical and phenomenological points of view. There is a lot of important results obtained within perturbation theory, which works well for high energy processes. On the other hand, the low energy hadron physics, in particular, the quark confinement, turns out to be dominated by nonperturbative effects, for which there is no rigorous theoretical explanation yet. To study nonperturbative aspects, a variety of different concepts and mathematical methods has been developed.

One of the main reasons, which makes a nonabelian gauge theory so different from other field theories is the rich mathematical structure of its classical configuration space (the gauge orbit space) and the corresponding phase space. The gauge orbit space is obtained by factorizing the infite-dimensional affine space of gauge potentials by the action of the group of local gauge transformations. Since this action is not free, non-generic orbit types (singularities) occur endowing the gauge orbit space with a stratified structure. Let us discuss some aspects indicating its physical relevance.

First, the geometry and topology of the generic (principal) stratum has been clarified a long time ago [33, 40]. In particular, one gets an intrinsic topological interpretation of the Gribov-ambiguity [18]. We stress that the problem of finding all Gribov copies has been discussed within specific models, see e.g. [28]. For a detailed analysis in the case of 2-dimensional cylindrical space time (including the Hamiltonian path integral) we refer to [39]. Investigating the topology of the determinant line bundle over the generic stratum, one gets an understanding of anomalies in terms of the family index theorem [1, 5], see also [11] for the Hamiltonian approach. In particular, one gets anomalies of purely topological type [47] which cannot be seen by perturbative quantum field theory.

There are partial results and conjectures concerning the relevance of nongeneric strata. First, generally speaking, nongeneric gauge orbits affect the classical motion on the orbit space due to boundary conditions and, in this way, they may produce nontrivial contributions to the path integral. They may lead to localization of certain quantum states, as it was suggested by finite-dimensional examples [14], see also [22] for a finite-dimensional gauge model. Further, the gauge field configurations belonging to nongeneric orbits can possess a magnetic charge, i.e. they can be considered as a kind of magnetic monopole configurations. Following t'Hooft [46], these could be responsible for quark confinement. The role of these configurations was investigated within the framework of Schrödinger quantum mechanics on the gauge orbit space of topological Chern-Simons theory in [2], see also [3] for an approach to 4-dimensional Yang-Mills theories with θ-term. Within t'Hooft's concept, the idea of abelian projection is of special importance and has been discussed by many authors. This concept was studied within the setting of quantum field theory at finite temperature on the 4-torus in [16, 17]. There, a hierarchy of defects, which should be related to the gauge orbit space structure, was discovered. Finally, let us also mention that in [21] the existence of additional anomalies corresponding to non-generic strata was suggested.

Most of the problems mentioned here are still awaiting a systematic investigation. For that purpose, a deeper insight into the structure of the gauge orbit space is necessary. The stratified structure of the full gauge orbit space was investigated in detail in [27]. Based on that, in a series of papers [34, 35, 37], see also [36] for a review, we have given a complete solution to the problem of determining the strata (including their partial ordering) for gauge theories with structure group $\mathrm{SU}(n)$ on closed connected manifolds of dimension $d = 2, 3, 4$. Our analysis was based in particular on the observation, made e.g. in [27], that orbit types are in bijective correspondence with a certain type of bundle reductions of the principal bundle underlying the gauge theory under consideration. We call bundle reductions of this type holonomy-induced Howe subbundles. This observation yields a method for solving the classification problem for orbit types, because it turns out that holonomy-induced Howe subbundles can be classified by methods of algebraic topology. In the present paper we extend the above results to the case of structure group $\mathrm{O}(n)$, $\mathrm{SO}(n)$ or $\mathrm{Sp}(n)$. We apply the same general method, but the classification problem for the subbundles turns out to be much harder than in the case of $\mathrm{SU}(n)$.

Let us put this paper into a broader perspective. In a next step, we will investigate the structure of the phase space (infinite-dimensional Hamiltonian system with symmetry) by applying the method of singular Marsden-Weistein reduction. This will yield the reduced phase space as a Hamiltonian system with singularities, which can be taken as a starting point for constructing the quantum theory. To separate the geometric problems arising from the reduction process from the analytic problems due to infinite dimensions, it is reasonable to study finite-dimensional approximations (provided by lattice gauge theory) of the above situation. We refer to [12, 13, 15, 22] for first results in this direction. In particular, in [22] we have studied the role of non-generic strata on the quantum level for one of these finite-dimensional models.

The paper is organized as follows: In Section 2 we present the method and formulate the program for solving the classification problem. In Section 3 we determine the Howe subgroups of O(n), SO(n) and Sp(n). In Section 4 we classify principal bundles whose structure group is a Howe subgroup under some assumptions concerning the space time manifold. This chapter is the heart of the paper. In Section 5 we determine the Howe subbundles of a given principal bundle and in Section 6 we specify the Howe subbundles which are holonomy-induced. Section 7 contains the factorization with respect to the action of the structure group on bundle reductions and in Section 8 we summarize the classification result. As an illustration, in Section 9 we apply the results to gauge theories with structure group O(4), SO(4) or Sp(2), defined over a space-time manifold diffeomorphic to $\mathbb{C}\mathrm{P}^2$ or $\mathrm{S}^2 \times \mathrm{S}^2$. For the convenience of the reader, the paper is supplemented by 5 Appendices, where we collect relevant material from algebraic topology referred to in the text.

Finally, we note that we restrict attention to closed (compact, without boundary) and simply connected space time manifolds of dimension 4. However, the topological results derived in Section 4 are obtained for CW-complexes, some of them hold true without the above assumptions.

2 Orbit types and holonomy-induced Howe subbundles

Let M be a closed connected orientable manifold, let G be a compact connected Lie-group and let P be a principal bundle over M with structure group G. We denote the affine Hilbert-space of connection forms on P of an appropriate Sobolev class k by $\mathcal{A}$ the Hilbert-Lie group of vertical automorphisms of P of Sobolev class $(k+1)$ by $\mathcal{G}$. For the analytic framework, see [33, 40]. The results we are going to derive are independent of the choice of k provided k is large enough.

Our aim is to investigate the structure of the configuration space of a gauge theory on P,

$$\mathcal{M} = \mathcal{A}/\mathcal{G}\,,$$

where the action of $\mathcal{G}$ on $\mathcal{A}$ is given by

$$(g, A) \mapsto A^{(g)} = g^{-1}Ag + g^{-1}\mathrm{d}g.$$

This space is known as the gauge orbit space. Since, in general, the action of $\mathcal{G}$ is not free, $\mathcal{M}$ is not a smooth manifold. However, the action is proper and admits slices [27]. Therefore, $\mathcal{M}$ is a stratified space, where the stratification is induced by the orbit types of the action of $\mathcal{G}$ on $\mathcal{A}$. Let us briefly recall the construction. The stabiliser (or isotropy group) of a connection $A \in \mathcal{A}$ is given by $\mathcal{G}_A = \{g \in \mathcal{G} \mid A^{(g)} = A\}$. It transforms under the action of $\mathcal{G}$ like $\mathcal{G}_{A^{(g)}} = g^{-1}\mathcal{G}_A g$. This allows to define the type of an orbit $[A]$ to be the conjugacy class of the subgroup $\mathcal{G}_A$ in $\mathcal{G}$. For given orbit type τ let $\mathcal{M}_\tau$ denote the subset of $\mathcal{M}$ of orbits of type τ. This defines a disjoint decomposition

$$\mathcal{M} = \bigcup_{\text{orbit types}} \mathcal{M}_\tau\,. \tag{1}$$

The following was shown in [27]:
- each $\mathcal{M}_\tau$ carries the structure of a Hilbert manifold,
- the orbit type deomposition (1) is locally finite,
- the frontier condition holds: if $\mathcal{M}_\tau \cap \overline{\mathcal{M}_\sigma} \neq \emptyset$ then $\mathcal{M}_\tau \subset \overline{\mathcal{M}_\sigma}$.

In [26, 27], the decomposition (1) is therefore called a stratification and the orbit type subsets $\mathcal{M}_\tau$ are referred to as the strata of $\mathcal{M}$. We adopt this terminology[1]. It was also shown in [27] that the natural partial ordering of orbit types, which is induced from the inclusion relation between subgroups of $\mathcal{G}$, satisfies

$$\tau \leq \sigma \quad \Leftrightarrow \quad \mathcal{M}_\sigma \subseteq \overline{\mathcal{M}_\tau}\,.$$

Thus, the the partially ordered set of orbit types contains information about which strata occur in the gauge theory on P and how they are glued together. In the present paper we determine the orbit types for structure groups $G = \mathrm{O}(n)$, $\mathrm{SO}(n)$ and $\mathrm{Sp}(n)$ and for M being a closed simply connected manifold of dimension 4; the partial ordering will be studied in a separate paper. For that purpose, we use a relation between orbit types and certain types of bundle reductions of P which for the first time was described in detail in [27]. Let us explain this relation. A subgroup $H \subseteq G$ with the property $C_G^2(H) = H$ is called *Howe subgroup*. For any subgroup H of G there exists a smallest Howe subgroup $\tilde{H}$ of G such that $H \subseteq \tilde{H}$. We say that $\tilde{H}$ is the Howe subgroup generated by H. $\tilde{H}$ can be obtained by taking the double centralizer, $\tilde{H} = \mathrm{C}_G^2(H)$. A reduction of P to a Howe subgroup is called *Howe subbundle*. For any reduction Q of P to a subgroup H there exists a smallest Howe subbundle $\tilde{Q}$ such that $Q \subseteq \tilde{Q}$. We say that $\tilde{Q}$ is the Howe subbundle generated by Q. $\tilde{Q}$ can be obtained from Q by extending the structure group to the Howe subgroup generated by H, i.e., $\tilde{Q} = Q \cdot \mathrm{C}_G(H)$. A Howe subbundle is called *holonomy-induced* if it is generated by the holonomy subbundle of some connection in P. Due to existence results for holonomy subbundles [25, Ch. II, Thm. 8.2] in dimension $d \geq 2$ this is equivalent to requiring that the Howe subbundle be generated by a connected reduction of P.

Theorem 2.1 (Kondracki and Rogulski [27]). *If* $\dim M > 1$, *there is an (order-preserving) bijection from the set of orbit types onto the set of isomorphism classes of holonomy-induced Howe subbundles, factorized by the action of the structure group* G *on bundle reductions.*

For a detailed proof, see [34]. The idea of the proof is based on the observation that when gauge transformations are viewed as equivariant maps from P to G, the stabilizer of a connection consists of the gauge transformations which are constant on any holonomy subbundle of that connection. It follows that the stabilizer is the same for any two connections whose holonomy subbundles (based at the same point) generate the same Howe subbundle. Therefore, stabilizers correspond to holonomy-induced Howe subbundles. Passing to classes on the level of the subbundles then corresponds to passing to conjugacy classes on the level of the stabilizers.

According to Theorem 2.1, in order to determine the orbit types of the action of $\mathcal{G}$ on $\mathcal{A}$ we have to do the following.

1. Determine the Howe subgroups of G.
2. Determine the Howe subbundles of P up to isomorphy.
3. Specify the Howe subbundles which are holonomy-induced.
4. Factorize by the action of the structure group G on bundle reductions.

To work out the full program for $G = \mathrm{O}(n)$, $\mathrm{SO}(n)$ or $\mathrm{Sp}(n)$ we have to assume that M is simply connected and of dimension 4. Step 1 will be treated in Section 3. Step 2 is the hardest one, it will be worked out in Sections 4 and 5. Steps 3 and 4 are treated in Sections 6 and 7, respectively. The results are summarized in Section 8.

---

[1] There exist several stronger notions of stratification.

As mentioned above, for structure group $G = \mathrm{SU}(n)$ the orbit types and their partial ordering have already been determined in [34, 35], see also [37]. Here the assumption that M be simply connected can be dropped. For the sake of completeness, we include these results into the summary in Section 8.

Remark 2.2. The assumption that $\dim(M)$ be exactly 4 is just made for simplicity. Our results on the classification of Howe subbundles hold also in dimension $d \leq 4$. In view of Theorem 2.1, we thus derive the orbit types for $\dim(M) = 2, 3, 4$.

3 Howe subgroups

Let $\mathbb{K} = \mathbb{R}, \mathbb{C}, \mathbb{H}$ denote the real numbers, complex numbers and quaternions, respectively. Let $\mathrm{I}_{\mathbb{K}}(n)$ denote the isometry groups of the standard scalar products on $\mathbb{K}^n$. The standard notation is obtained by replacing $\mathrm{I}_{\mathbb{R}} = \mathrm{O}$, $\mathrm{I}_{\mathbb{C}} = \mathrm{U}$ and $\mathrm{I}_{\mathbb{H}} = \mathrm{Sp}$. For a group G and a subgroup $H \subseteq G$, let $\mathrm{C}_G(H)$ and $\mathrm{N}_G(H)$ denote the centralizer and the normalizer, respectively, of H in G.

The study of the Howe subgroups of $\mathrm{O}(n)$ and $\mathrm{Sp}(n)$ amounts to an application of the theory of finite-dimensional real and quaternionic von Neumann algebras, respectively. We cite the results from [38] where the more general case of the isometry group of a Hermitian form over $\mathbb{K}$ was treated. The Howe subgroups of $\mathrm{SO}(n)$ can be derived from those of $\mathrm{O}(n)$. Since to our knowledge this is not documented in the literature, we give full proofs here. For this discussion, and also for the discussion of the property of a bundle reduction to be holonomy-induced in Section 6, we need information about the inclusion relations between Howe subgroups of $\mathrm{O}(n)$ and $\mathrm{Sp}(n)$. Therefore, for these groups, from [38] we also cite the operations producing the direct predecessors (up to conjugacy) of a given Howe subgroup w.r.t. the natural partial ordering defined by the inclusion relation.

Let $G = \mathrm{I}_{\mathbb{K}}(n)$ be given. A Howe subgroup H of G is called irreducible if the representation of $H \times \mathrm{C}_G(H)$ on $\mathbb{K}^n$ is irreducible. Orthogonally decomposing $\mathbb{K}^n = \mathbb{K}^{n_1} \oplus \cdots \oplus \mathbb{K}^{n_r}$ into $H \times \mathrm{C}_G(H)$-irreducible subspaces one obtains a decomposition of H into a direct product of irreducible Howe subgroups of $\mathrm{I}_{\mathbb{K}}(n_i)$. The centralizer of H is then the direct product of the centralizers of the irreducible factors of H in the groups $\mathrm{I}(n_i)$.

3.1 Howe subgroups of $\mathrm{O}(n)$

For any positive integer m, field restriction defines homomorphisms

$$\varphi_{\mathbb{C},\mathbb{R}} : \mathrm{M}_{\mathbb{C}}(m) \to \mathrm{M}_{\mathbb{R}}(2m)\,, \qquad \varphi_{\mathbb{H},\mathbb{C}} : \mathrm{M}_{\mathbb{H}}(m) \to \mathrm{M}_{\mathbb{C}}(2m)\,, \qquad \varphi_{\mathbb{H},\mathbb{R}} : \mathrm{M}_{\mathbb{H}}(m) \to \mathrm{M}_{\mathbb{R}}(4m)\,.$$

For convenience, we set $\varphi_{\mathbb{R},\mathbb{R}} = \mathrm{id}_{\mathrm{M}_{\mathbb{R}}(m)}$. Explicitly, we choose

$$\varphi_{\mathbb{C},\mathbb{R}}(a_1 + \mathrm{i}a_2) = \begin{bmatrix} a_1 & a_2 \\ -a_2 & a_1 \end{bmatrix}, \qquad a_k \in \mathrm{M}_{\mathbb{R}}(m)\,, \tag{2}$$

$$\varphi_{\mathbb{H},\mathbb{C}}(a_1 + a_2\mathrm{j}) = \begin{bmatrix} \overline{a_1} & -\overline{a_2} \\ a_2 & a_1 \end{bmatrix}, \qquad a_k \in \mathrm{M}_{\mathbb{C}}(m)\,, \tag{3}$$

and $\varphi_{\mathbb{H},\mathbb{R}} := \varphi_{\mathbb{C},\mathbb{R}} \circ \varphi_{\mathbb{H},\mathbb{C}}$. I.e.,

$$\varphi_{\mathbb{H},\mathbb{R}}(a_1 + \mathrm{i}a_2 + \mathrm{j}a_3 + \mathrm{k}a_4) = \begin{bmatrix} a_1 & -a_2 & -a_3 & -a_4 \\ a_2 & a_1 & a_4 & -a_3 \\ a_3 & -a_4 & a_1 & a_2 \\ a_4 & a_3 & -a_2 & a_1 \end{bmatrix}, \qquad a_k \in \mathrm{M}_{\mathbb{R}}(m)\,. \tag{4}$$

These mappings commute with taking adjoints w.r.t. the respective scalar product. Therefore, they induce Lie group embeddings (denoted by the same symbol)

$$\varphi_{\mathbb{C},\mathbb{R}} : \mathrm{U}(m) \to \mathrm{O}(2m)\,, \qquad \varphi_{\mathbb{H},\mathbb{C}} : \mathrm{Sp}(m) \to \mathrm{U}(2m)\,, \qquad \varphi_{\mathbb{H},\mathbb{R}} : \mathrm{Sp}(m) \to \mathrm{O}(4m)\,.$$

Since $\mathrm{U}(m)$ is connected, $\varphi_{\mathbb{C},\mathbb{R}}(\mathrm{U}(m)) \subseteq \mathrm{SO}(2m)$ and hence $\varphi_{\mathbb{H},\mathbb{R}}(\mathrm{Sp}(n)) \subseteq \mathrm{SO}(4m)$. Moreover, $\varphi_{\mathbb{H},\mathbb{C}}(\mathrm{Sp}(n)) \subseteq \mathrm{SU}(2m)$.

The conjugacy classes of irreducible Howe subgroups of $\mathrm{O}(n_i)$ are in bijective correspondence with the solutions of the equation $\delta_i m_i k_i = n_i$, where (m_i, k_i) is an ordered pair of positive integers and $\delta_i = 1, 2, 4$. For given δ_i, m_i, k_i, a representative for the corresponding class of Howe subgroups is given by $H_i = \mathrm{I}_{\mathbb{K}_i}{}^{k_i}_{m_i}$ where

$$\mathrm{I}_{\mathbb{K}_i}{}^{k_i}_{m_i} = \{\varphi_{\mathbb{K},\mathbb{R}}(a) \otimes_{\mathbb{R}} \mathbb{1}_{k_i} \;:\; a \in \mathrm{I}_{\mathbb{K}_i}(m_i)\} \tag{5}$$

w.r.t. the decomposition $\mathbb{R}^{n_i} = \mathbb{R}^{\delta_i m_i} \otimes_{\mathbb{R}} \mathbb{R}^{k_i}$ or, alternatively,

$$\mathrm{I}_{\mathbb{K}_i}{}^{k_i}_{m_i} = \left\{\varphi_{\mathbb{K},\mathbb{R}}(a) \oplus \overset{k_i}{\cdots} \oplus \varphi_{\mathbb{K},\mathbb{R}}(a) \;:\; a \in \mathrm{I}_{\mathbb{K}_i}(m_i)\right\} \tag{6}$$

w.r.t. the decomposition $\mathbb{R}^{n_i} = \mathbb{R}^{\delta_i m_i} \oplus \overset{k_i}{\cdots} \oplus \mathbb{R}^{\delta_i m_i}$. Here $\mathbb{K}_i = \mathbb{R}$ for $\delta_i = 1$, $\mathbb{K}_i = \mathbb{C}$ for $\delta_i = 2$ and $\mathbb{K}_i = \mathbb{H}$ for $\delta_i = 4$. We call $\mathbb{K}_i$ the base field, m_i the rank and k_i the multiplicity of H_i. Moreover, we denote $\mathrm{O}^{k_i}_{m_i} = \mathrm{I}_{\mathbb{R}}{}^{k_i}_{m_i}$, $\mathrm{U}^{k_i}_{m_i} = \mathrm{I}_{\mathbb{C}}{}^{k_i}_{m_i}$ and $\mathrm{Sp}^{k_i}_{m_i} = \mathrm{I}_{\mathbb{H}}{}^{k_i}_{m_i}$. A representative for the conjugacy class of the centralizer of H_i is given by $\mathrm{I}_{\mathbb{K}_i}{}^{m_i}_{k_i}$ (i.e., rank and multiplicity are interchanged). Thus, up to conjugacy, the Howe subgroups of $\mathrm{O}(n)$ are given by the direct products

$$H = \mathrm{I}_{\mathbb{K}_1}{}^{k_1}_{m_1} \times \cdots \times \mathrm{I}_{\mathbb{K}_r}{}^{k_r}_{m_r}\,, \qquad \sum\nolimits_{i=1}^{r} \dim_{\mathbb{R}} \mathbb{K}_i\, m_i k_i = n\,, \qquad r = 1, 2, 3, \dots\,. \tag{7}$$

where the factors are embedded by virtue of the homomorphisms $\varphi_{\mathbb{K}_i,\mathbb{R}}$. The factors are referred to as O-factors, U-factors and Sp-factors according to the base field being $\mathbb{R}$, $\mathbb{C}$ or $\mathbb{H}$, respectively. A representative for the conjugacy class of the centralizer of H is obtained from H by interchanging ranks and multiplicities. The subgroup $\mathrm{I}_{\mathbb{R}}{}^{n}_{1} \equiv \mathrm{O}^{n}_{1}$ coincides with the center of $\mathrm{O}(n)$ and the subgroup $\mathrm{I}_{\mathbb{R}}{}^{1}_{n} \equiv \mathrm{O}^{1}_{n}$ coincides with $\mathrm{O}(n)$ itself.

The operations which produce the direct predecessors of a given Howe subgroup of $\mathrm{O}(n)$ (up to conjugacy) are

– *Merging:* replacing a double factor $\mathrm{I}_{\mathbb{K}}{}^{k_1}_{m} \times \mathrm{I}_{\mathbb{K}}{}^{k_2}_{m}$ by a single factor $\mathrm{I}_{\mathbb{K}}{}^{k_1+k_2}_{m}$; $\mathrm{I}_{\mathbb{K}}(m)$ is diagonally embedded into $\mathrm{I}_{\mathbb{K}}(m) \times \mathrm{I}_{\mathbb{K}}(m)$;

– *Splitting:* replacing a factor $\mathrm{I}_{\mathbb{K}}{}^{k}_{m}$ with $m > 1$ by the double factor $\mathrm{I}_{\mathbb{K}}{}^{k}_{m_1} \times \mathrm{I}_{\mathbb{K}}{}^{k}_{m_2}$, where $m_1 + m_2 = m$; $\mathrm{I}_{\mathbb{K}}(m_1) \times \mathrm{I}_{\mathbb{K}}(m_2)$ is embedded into $\mathrm{I}_{\mathbb{K}}(m)$ in the obvious way;

– *Inverse field restriction:* replacing a factor O^{k}_{2m} or U^{k}_{2m} by a factor U^{k}_{m} or Sp^{k}_{m}, embedded via $\varphi_{\mathbb{C},\mathbb{R}}$ or $\varphi_{\mathbb{H},\mathbb{C}}$, respectively;

– *Inverse field extension:* replacing a factor U^{k}_{m} or Sp^{k}_{m} by a factor O^{2k}_{m} or U^{2k}_{m}, embedded in the obvious way.

As an example, the Hasse diagrams of the sets of conjugacy classes of Howe subgroups of $\mathrm{O}(2)$, $\mathrm{O}(3)$ and $\mathrm{O}(4)$ are displayed in Figure 1.

We will also have to study the identity connected component H_0 of a Howe subgroup H. One has

$$H_0 = (\mathrm{I}_{\mathbb{K}_1}{}^{k_1}_{m_1})_0 \times \cdots \times (\mathrm{I}_{\mathbb{K}_r}{}^{k_r}_{m_r})_0\,,$$

where $(\mathrm{I}_{\mathbb{K}_i}{}^{k_i}_{m_i})_0$ denotes the identity connected component of the factor $\mathrm{I}_{\mathbb{K}_i}{}^{k_i}_{m_i}$. For $\mathbb{K}_i = \mathbb{R}$, this is $(\mathrm{I}_{\mathbb{R}}{}^{k_i}_{m_i})_0 \equiv \mathrm{SO}^{k_i}_{m_i}$, whereas for $\mathbb{K}_i = \mathbb{C}, \mathbb{H}$, it coincides with $\mathrm{I}_{\mathbb{K}_i}{}^{k_i}_{m_i}$. Thus, H_0 consists of SO-factors, U-factors and Sp-factors.


$\mathrm{O}_1^1\,\mathrm{O}_1^1$
O_1^2
U_1^1
O_2^1
$\mathrm{O}_1^1\,\mathrm{O}_1^1\,\mathrm{O}_1^1$
O_1^3
$\mathrm{O}_1^1\,\mathrm{O}_1^2$
$\mathrm{O}_1^1\,\mathrm{U}_1^1$
$\mathrm{O}_1^1\,\mathrm{O}_2^1$
O_3^1
$\mathrm{O}_1^1\,\mathrm{O}_1^1\,\mathrm{O}_1^1\,\mathrm{O}_1^1$
$\mathrm{O}_1^1\,\mathrm{O}_1^3$
$\mathrm{O}_1^1\,\mathrm{O}_1^1\,\mathrm{O}_1^2$
$\mathrm{O}_1^2\,\mathrm{O}_2^1$
$\mathrm{O}_1^1\,\mathrm{O}_1^1\,\mathrm{O}_2^1$
$\mathrm{O}_1^1\,\mathrm{O}_3^1$
$\mathrm{O}_1^2\,\mathrm{O}_1^2$
O_1^4
O_4^1
$\mathrm{O}_1^2\,\mathrm{U}_1^1$
$\mathrm{O}_1^1\,\mathrm{O}_1^1\,\mathrm{U}_1^1$
$\mathrm{O}_2^1\,\mathrm{U}_1^1$
$\mathrm{O}_2^1\,\mathrm{O}_2^1$
$\mathrm{U}_1^1\,\mathrm{U}_1^1$
U_1^2
O_2^2
U_2^1
Sp_1^1

Figure 1: Hasse diagrams of the sets of conjugacy classes of Howe subgroups of O(2), O(3), O(4). For brevity, direct product signs are omitted, i.e., $\mathrm{O}_1^1\,\mathrm{O}_1^1$ stands for $\mathrm{O}_1^1 \times \mathrm{O}_1^1$ etc.

3.2 Howe subgroups of $\mathrm{Sp}(n)$

We use the convention that on $\mathbb{H}^n$, scalars act by multiplication from the right and endomorphisms by matrix multiplication from the left. The conjugacy classes of irreducible Howe subgroups of $\mathrm{Sp}(n_i)$ are in bijective correspondence with ordered pairs of positive integers (m_i, k_i) such that $m_i k_i = n_i$, together with a choice of $\mathbb{K}_i = \mathbb{R}, \mathbb{C}, \mathbb{H}$. For given (m_i, k_i) and $\mathbb{K}_i$, a representative for the corresponding class of Howe subgroups is given by $H_i = \mathrm{I}_{\mathbb{K}_i}{}^{k_i}_{m_i}$, where

$$\mathrm{I}_{\mathbb{K}_i}{}^{k_i}_{m_i} = \{a \otimes_{\mathbb{H}} \mathbb{1}_{k_i} \ : \ a \in \mathrm{I}_{\mathbb{K}_i}(m_i)\} \tag{8}$$

w.r.t. the decomposition $\mathbb{H}^{n_i} = \mathbb{H}^{m_i} \otimes_{\mathbb{H}} \mathbb{H}^{k_i}$ or, alternatively,

$$\mathrm{I}_{\mathbb{K}_i}{}^{k_i}_{m_i} = \left\{ a \oplus \overset{k_i}{\cdots} \oplus a \ : \ a \in \mathrm{I}_{\mathbb{K}_i}(m_i) \right\} \tag{9}$$

w.r.t. the decomposition $\mathbb{H}^{n_i} = \mathbb{H}^{m_i} \oplus \overset{k_i}{\cdots} \oplus \mathbb{H}^{m_i}$. Recall that the tensor product over $\mathbb{H}$ is defined w.r.t. the actions of $\mathbb{H}$ by right multiplication on the first factor and by left multiplication on the second factor. Thus, operators of the type $a \otimes_{\mathbb{H}} b$ can be defined provided the entries of b are central, i.e., real. We will use the same terminology as in the case of $\mathrm{O}(n)$, i.e., $\mathbb{K}_i$ will be referred to as the base field of H_i, m_i as the rank of H_i and k_i as the multiplicity of H_i and we will speak of O-, U- and Sp-factors. A representative for the conjugacy class of the centralizer of $\mathrm{I}_{\mathbb{K}_i}{}^{k_i}_{m_i}$ is given by $\mathrm{I}_{\mathbb{L}_i}{}^{m_i}_{k_i}$, where $\mathbb{L}_i = \mathbb{H}$ if $\mathbb{K}_i = \mathbb{R}$, $\mathbb{L}_i = \mathbb{R}$ if $\mathbb{K}_i = \mathbb{H}$ and $\mathbb{L}_i = \mathbb{C}$ otherwise.
Thus, up to conjugacy, the Howe subgroups of $\mathrm{Sp}(n)$ are given by direct products

$$H = \mathrm{I}_{\mathbb{K}_1}{}^{k_1}_{m_1} \times \cdots \times \mathrm{I}_{\mathbb{K}_r}{}^{k_r}_{m_r}\,, \qquad n = \sum\nolimits_{i=1}^{r} m_i k_i\,,\,, \qquad r = 1, 2, 3, \ldots\,, \tag{10}$$


O_1^1
U_1^1
Sp_1^1
O_2^1
$\mathrm{O}_1^1\,\mathrm{Sp}_1^1$
Sp_1^2
$\mathrm{O}_1^1\,\mathrm{O}_1^1$
$\mathrm{O}_1^1\,\mathrm{U}_1^1$
$\mathrm{U}_1^1\,\mathrm{Sp}_1^1$
$\mathrm{Sp}_1^1\,\mathrm{Sp}_1^1$
O_1^2
Sp_2^1
U_1^2
$\mathrm{U}_1^1\,\mathrm{U}_1^1$
U_2^1

Figure 2: Hasse diagrams of the sets of conjugacy classes of Howe subgroups for Sp(1) and Sp(2). For the notation, see Figure 1. For Sp(1) one has the identifications $\mathrm{Sp}_1^1 = \mathrm{Sp}(1) \equiv \mathrm{SU}(2)$, $\mathrm{U}_1^1 =$ maximal toral subgroup of SU(2) and $\mathrm{O}_1^1 \equiv \mathbb{Z}_2 =$ center of SU(2).

and a representative for the conjugacy class of the centralizer of H is obtained from H by interchanging ranks and multiplicities as well as base fields $\mathbb{R}$ and $\mathbb{H}$. The subgroup $\mathrm{I}_{\mathbb{R}}{}^{n}_{1} \equiv \mathrm{O}_1^n$ coincides with the center of $\mathrm{Sp}(n)$ and the subgroup $\mathrm{I}_{\mathbb{H}}{}^{1}_{n} \equiv \mathrm{Sp}_n^1$ coincides with $\mathrm{Sp}(n)$ itself.
The operations which produce the direct predecessors of a given Howe subgroup of $\mathrm{Sp}(n)$ (up to conjugacy) are

– *Merging* and *splitting* similar to the case of $\mathrm{O}(n)$
– *Inverse field restriction:* replacing a factor O_{2m}^k or U_{2m}^k by a factor U_m^{2k} or Sp_m^{2k}, embedded via $\varphi_{\mathbb{C},\mathbb{R}}$ or $\varphi_{\mathbb{H},\mathbb{C}}$, respectively;
– *Inverse field extension:* replacing a factor U_m^k or Sp_m^k by a factor O_m^k or U_m^k, respectively, embedded in the obvious way.

3.3 Howe subgroups of $\mathrm{SO}(n)$

First, we treat the case of odd n. Here

$$\mathrm{O}(n) = Z \cdot \mathrm{SO}(n)\,, \tag{11}$$

where $Z = \{\mathbb{1}, -\mathbb{1}\}$ is the center of $\mathrm{O}(n)$.

Proposition 3.1. *For n odd, intersection with* $\mathrm{SO}(n)$ *defines a bijection from the set of Howe subgroups of* $\mathrm{O}(n)$ *onto the set of Howe subgroups of* $\mathrm{SO}(n)$*. The bijection preserves the equivalence relation of conjugacy. Its inverse is given by multiplication by the center Z.*

Proof. We start with deriving some formulae. As an immediate consequence of (11), for any subgroup H of $\mathrm{O}(n)$,

$$\mathrm{C}_{\mathrm{O}(n)}(H) = \mathrm{C}_{\mathrm{O}(n)}(Z \cdot H)\,. \tag{12}$$

Furthermore, the chain of inclusions $\mathrm{SO}(n) \cap H \subseteq H \subseteq Z \cdot (\mathrm{SO}(n) \cap H)$ implies

$$\mathrm{C}_{\mathrm{O}(n)}(\mathrm{SO}(n) \cap H) \supseteq \mathrm{C}_{\mathrm{O}(n)}(H) \supseteq \mathrm{C}_{\mathrm{O}(n)}\big(Z \cdot (\mathrm{SO}(n) \cap H)\big)\,.$$

Then (12), applied to the subgroup $\mathrm{SO}(n) \cap H$, yields

$$\mathrm{C}_{\mathrm{O}(n)}(H) = \mathrm{C}_{\mathrm{O}(n)}(\mathrm{SO}(n) \cap H)\,. \tag{13}$$

Finally, by writing the centralizer in $\mathrm{SO}(n)$ as $\mathrm{C}_{\mathrm{SO}(n)} = \mathrm{SO}(n) \cap \mathrm{C}_{\mathrm{O}(n)}$ one sees that (11) implies that, for any subgroup $\tilde{H}$ of $\mathrm{SO}(n)$,

$$Z \cdot \mathrm{C}_{\mathrm{SO}(n)}(\tilde{H}) = \mathrm{C}_{\mathrm{O}(n)}(\tilde{H}) . \tag{14}$$

Now let H be a Howe subgroup of $\mathrm{O}(n)$. Then $H = \mathrm{C}_{\mathrm{O}(n)}(H')$ for some subgroup H' of $\mathrm{O}(n)$. According to (13), then $H = \mathrm{C}_{\mathrm{O}(n)}\big(\mathrm{SO}(n) \cap H'\big)$. Intersection with $\mathrm{SO}(n)$ yields

$$\mathrm{SO}(n) \cap H = \mathrm{C}_{\mathrm{SO}(n)}\big(\mathrm{SO}(n) \cap H'\big) .$$

Hence, $\mathrm{SO}(n) \cap H$ is a Howe subgroup of $\mathrm{SO}(n)$. Conversely, let $\tilde{H}$ be a Howe subgroup of $\mathrm{SO}(n)$. Consider $H = Z \cdot \tilde{H}$. By construction, $\tilde{H} = \mathrm{SO}(n) \cap H$. Moreover, since $\tilde{H} = \mathrm{C}_{\mathrm{SO}(n)}(\tilde{H}')$ for some subgroup $\tilde{H}'$ of $\mathrm{SO}(n)$, (14) implies $H = \mathrm{C}_{\mathrm{O}(n)}(\tilde{H}')$. Hence, H is a Howe subgroup of $\mathrm{O}(n)$.
Finally, in view of (11) it is obvious that subgroups of $\mathrm{O}(n)$ that are conjugate under $\mathrm{O}(n)$ are also conjugate under $\mathrm{SO}(n)$. ☐

When n is even, there exist Howe subgroups of $\mathrm{O}(n)$ whose intersection with $\mathrm{SO}(n)$ is not a Howe subgroup of $\mathrm{SO}(n)$. The simplest example of such a situation is provided by the center of $\mathrm{O}(2)$ which is contained in $\mathrm{SO}(2)$ but is not a Howe subgroup there, because $\mathrm{SO}(2)$ is Abelian. Moreover, distinct Howe subgroups of $\mathrm{O}(n)$ may have the same intersection with $\mathrm{SO}(n)$. Again, the simplest example is provided by $\mathrm{O}(2)$, and the Howe subgroups $\mathrm{SO}(2)$ and $\mathrm{O}(2)$.
To begin with, let us introduce some notation. For a subgroup H of $\mathrm{O}(n)$ let

$$\mathrm{C}H := \mathrm{C}_{\mathrm{O}(n)}(H) , \qquad \mathrm{S}H := \mathrm{SO}(n) \cap H , \qquad \mathrm{M}H := \mathrm{C}^2\mathrm{S}H .$$

By construction, $\mathrm{M}H$ is the Howe subgroup of $\mathrm{O}(n)$ generated by $\mathrm{S}H$, i.e., the smallest Howe subgroup of $\mathrm{O}(n)$ containing $\mathrm{S}H$. For convenience, C, S and M will be viewed as maps on the set of subgroups of $\mathrm{O}(n)$. The basic properties of these maps are monotony,

$$H \subseteq K \quad \Rightarrow \quad \mathrm{C}H \supseteq \mathrm{C}K , \quad \mathrm{C}^2H \subseteq \mathrm{C}^2K , \quad \mathrm{S}H \subseteq \mathrm{S}K , \quad \mathrm{M}H \subseteq \mathrm{M}K$$

and periodicity resp. idempotence,

$$\mathrm{C}^3 = \mathrm{C} , \qquad \mathrm{S}^2 = \mathrm{S} , \qquad \mathrm{M}^2 = \mathrm{M} .$$

For C, these properties are well known. For S, they are obvious. For M, monotony follows from that of C^2 and S. Idempotence can be seen as follows. $\mathrm{M}H$ is a Howe subgroup of $\mathrm{O}(n)$ containing $\mathrm{SM}H$. Since M^2H is the smallest such subgroup, $\mathrm{M}^2H \subseteq \mathrm{M}H$. Conversely, we have both $\mathrm{S}H \subset \mathrm{M}H$ and $\mathrm{SM}H \subset \mathrm{M}^2H$. Applying S to the first inclusion and composing with the second one we obtain $\mathrm{S}H \subseteq \mathrm{SM}H \subseteq \mathrm{M}^2H$. I.e., M^2H is a Howe subgroup containing $\mathrm{S}H$. As $\mathrm{M}H$ is the smallest such subgroup, $\mathrm{M}H \subseteq \mathrm{M}^2H$.
A Howe subgroup H of $\mathrm{O}(n)$ will be called S-admissible if $\mathrm{S}H$ is a Howe subgroup of $\mathrm{SO}(n)$ and there is no smaller Howe subgroup of $\mathrm{O}(n)$ containing $\mathrm{S}H$. Using M, this can be reformulated as follows. H is S-admissible if and only if $\mathrm{S}H$ is a Howe subgroup of $\mathrm{SO}(n)$ and $\mathrm{M}H = H$.

Proposition 3.2. S *induces a bijection from the set of* S*-admissible Howe subgroups of* $\mathrm{O}(n)$ *onto the set of Howe subgroups of* $\mathrm{SO}(n)$*. The inverse is given by* M.

Proof. By definition of S-admissibility, S defines a map from the set of S-admissible Howe subgroups of $\mathrm{O}(n)$ to the set of Howe subgroups of $\mathrm{SO}(n)$. Since the intersection of arbitrarily many Howe subgroups is a Howe subgroup, this map is injective. To see that it is also surjective, let $\tilde{H}$ be a Howe subgroup of $\mathrm{SO}(n)$. We show that $\tilde{H} = \mathrm{SM}\tilde{H}$ (then evidently $\mathrm{M}\tilde{H}$ is S-admissible and M is inverse to S). Since $\mathrm{S}\tilde{H} = \tilde{H}$, $\tilde{H} \subseteq \mathrm{M}\tilde{H}$. Application of S yields $\tilde{H} \subseteq \mathrm{SM}\tilde{H}$. Conversely, by $\mathrm{SC}\tilde{H} \subseteq \mathrm{C}\tilde{H}$ and monotony of C and S,

$$\mathrm{SM}\tilde{H} = \mathrm{SC}\big(\mathrm{C}\tilde{H}\big) \subseteq \mathrm{SC}\big(\mathrm{SC}\tilde{H}\big)\,.$$

The rhs. equals $\mathrm{C}^2_{\mathrm{SO}(n)}(\tilde{H})$. Since $\tilde{H}$ is a Howe subgroup of $\mathrm{SO}(n)$, $\mathrm{C}^2_{\mathrm{SO}(n)}(\tilde{H}) = \tilde{H}$ and hence $\mathrm{SM}\tilde{H} \subseteq \tilde{H}$. $\square$

Note that S-admissibility is not a necessary condition for a Howe subgroup of $\mathrm{O}(n)$ to yield a Howe subgroup of $\mathrm{SO}(n)$ by intersection. In fact, for Howe subgroups H that are not S-admissible, $\mathrm{S}H$ may or may not be a Howe subgroup of $\mathrm{SO}(n)$, as is shown by the Howe subgroups $H = \mathrm{O}(2)$ and $H =$ center of $\mathrm{O}(2)$.

Next, we determine the S-admissible Howe subgroups of $\mathrm{O}(n)$. As noted above, any such subgroup is stable under M.

Lemma 3.3. *Let H be an* M*-stable Howe subgroup of* $\mathrm{O}(n)$*. Then* $\mathrm{S}H$ *is a Howe subgroup of* $\mathrm{SO}(n)$ *if and only if* $H = (\mathrm{MC})^2 H$*.*

Proof. First, assume that $H = (\mathrm{MC})^2 H$ holds. Application of S yields $\mathrm{S}H = \mathrm{S}(\mathrm{MC})^2 H$. We observe: if K is a Howe subgroup of $\mathrm{O}(n)$ then $\mathrm{S}K \subseteq \mathrm{M}K \subseteq K$. Application of S yields $\mathrm{SM}K = \mathrm{S}K$. In particular, this implies $\mathrm{SMC} = \mathrm{SC}$. Using in addition the obvious relation $\mathrm{CM} = \mathrm{CS}$, the expression $\mathrm{S}(\mathrm{MC})^2 H$ can be rewritten as $\mathrm{SC}(\mathrm{SC}H)$, i.e., as the centralizer of the subgroup $\mathrm{SC}H$ of $SO(n)$, taken in $\mathrm{SO}(n)$. It follows that $\mathrm{S}H$ is a Howe subgroup of $\mathrm{SO}(n)$.
Conversely, assume that $\mathrm{S}H$ is a Howe subgroup of $\mathrm{SO}(n)$. We rewrite $(\mathrm{MC})^2 H$ as $\mathrm{C}^2(\mathrm{SC})^2 H$. Applying C to $H = \mathrm{M}H$ and using $\mathrm{CM} = \mathrm{CS}$ again we obtain $\mathrm{C}H = \mathrm{CS}H$. Hence, in the expression $\mathrm{C}^2(\mathrm{SC})^2 H$ we can replace H by $\mathrm{S}H$, thus obtaining $\mathrm{C}^2(\mathrm{SC})^2\mathrm{S}H$. Now $(\mathrm{SC})^2\mathrm{S}H$ is the double centralizer of $\mathrm{S}H$ in $\mathrm{SO}(n)$ and hence equals $\mathrm{S}H$. Upon using $\mathrm{C}H = \mathrm{CS}H$ once more, we arrive at $(\mathrm{MC})^2 H = H$, as asserted. $\square$

As a result, in order to determine the Howe subgroups of $\mathrm{SO}(n)$, we may first determine the Howe subgroups that are stable under M and then, among these, the Howe subgroups that are stable under $(\mathrm{MC})^2$.

Lemma 3.4. *A Howe subgroup H of* $\mathrm{O}(n)$ *is stable under* M *except for the following cases.*
(A) *H has a factor O_2^k with k odd and no other* O*-factor of odd multiplicity. Here,* $\mathrm{M}H$ *arises from H by inverse field restriction, i.e., by replacing O_2^k by U_1^k.*
(B) *H has a double factor $\mathrm{O}_1^k \times \mathrm{O}_1^l$ with k, l odd and no other* O*-factor of odd multiplicity. Here,* $\mathrm{M}H$ *arises from H by merging this double factor to O_1^{k+l}.*

Proof. First, we determine $\mathrm{M}H$ in cases (A) and (B). In case (A), let K be the Howe subgroup obtained from H by inverse field restriction of the factor O_2^k under consideration. Thus, this factor is replaced by the factor U_1^k. Since the original factor O_2^k is the only O-factor of odd multiplicity of H, an element of H has negative determinant if and only if its entry in this factor has so. Hence, $\mathrm{S}H = K$. Since K is Howe, then $\mathrm{M}H = K$. In case (B), let K be the Howe subgroup obtained from H by merging the double factor $\mathrm{O}_1^k \times \mathrm{O}_1^l$ to O_1^{k+l}. Since H has no other

O-factors of odd multiplicity, an element of H has negative determinant if and only if its entries in the two factors O_1^k and O_1^l are distinct. Hence, $\mathrm{S}H = K$ and so $\mathrm{M}H = K$. Thus, in cases (A) and (B), H is not stable under M.

Conversely, assume that H is not stable under M. Since $\mathrm{S}H$ has the same dimension as H, $\mathrm{M}H$ has the same dimension as H. It follows that H has a direct predecessor K, wrt. the natural partial ordering of Howe subgroups modulo conjugacy, of the same dimension. By comparing the dimension of the factors that are replaced by one another through the operations of merging, splitting, inverse field restriction and inverse field extension, one finds that there are only two situations where the dimension does not change. These are

- inverse field restriction of a factor O_2^k, which yields the factor U_1^k instead,
- merging of a double factor $\mathrm{O}_1^k \times \mathrm{O}_1^l$, which yields the factor O_1^{k+l}.

If, in the first situation, k is even or if H contains further O-factors of odd multiplicity, $\mathrm{S}H$ contains elements whose entry in the factor O_2^k under consideration has negative determinant, hence $\mathrm{S}K \neq \mathrm{S}H$. Similarly, in the second situation, if k or l is even or if H contains further O-factors of odd multiplicity, $\mathrm{S}H$ contains elements whose entries in the factors O_1^k and O_1^l are distinct. Thus, we are either in case (A) or case (B). $\square$

Lemma 3.5. *Let H be a Howe subgroup of $\mathrm{O}(n)$ which is stable under* M. *Then H is stable under* $(\mathrm{MC})^2$ *if and only if $\mathrm{C}H$ is stable under M.*

Proof. If $\mathrm{C}H$ is M-stable then $(\mathrm{MC})^2 H = \mathrm{MCMC}H = \mathrm{MC}^2 H = \mathrm{M}H = H$. If $\mathrm{C}H$ is not M-stable then by Lemma 3.4 $\mathrm{C}H$ and $\mathrm{MC}H$ are of the form

$$\mathrm{C}H = (\mathrm{C}H)^{(0)} \times \mathrm{O}_2^k\,, \qquad \mathrm{MC}H = (\mathrm{C}H)^{(0)} \times \mathrm{U}_1^k$$

or

$$\mathrm{C}H = (\mathrm{C}H)^{(0)} \times \mathrm{O}_1^k \times \mathrm{O}_1^l\,, \qquad \mathrm{MC}H = (\mathrm{C}H)^{(0)} \times \mathrm{O}_1^{k+l}\,,$$

where k and l are odd and $(\mathrm{C}H)^{(0)}$ does not contain an O-factor of odd multiplicity. Then H and $\mathrm{CMC}H$ are of the form

$$H = H^{(0)} \times \mathrm{O}_k^2\,, \qquad \mathrm{CMC}H = H^{(0)} \times \mathrm{U}_k^1$$

or

$$H = H^{(0)} \times \mathrm{O}(k)_1 \times \mathrm{O}_l^1\,, \qquad \mathrm{CMC}H = H^{(0)} \times \mathrm{O}_{k+l}^1\,,$$

respectively, where $H^{(0)}$ corresponds to $(\mathrm{C}H)^{(0)}$ under taking the centralizer. In both cases, H and $\mathrm{CMC}H$ have different dimension. On the other hand, $(\mathrm{MC})^2 H$ arises from $\mathrm{CMC}H$ by application of M and hence has the same dimension as $\mathrm{CMC}H$. Therefore, $H \neq (\mathrm{MC})^2 H$. $\square$

Lemmas 3.3–3.5 imply

Proposition 3.6. *A Howe subgroup of* $\mathrm{O}(n)$, *n even, is* S*-admissible if and only if neither itself nor its centralizer in* $\mathrm{O}(n)$ *belong to cases* (A) *or* (B) *of Lemma* 3.4. $\square$

For convenience, we reformulate the conditions for $\mathrm{C}H$ to belong to cases (A) or (B) of Lemma 3.4 as conditions on H. $\mathrm{C}H$ belongs to case (A) of Lemma 3.4 iff H has an O-factor of odd rank and multiplicity 2 and no further O-factor of odd rank. $\mathrm{C}H$ belongs to case (B) of Lemma 3.4 iff H has two O-factors of odd rank and multiplicity 1 and no further O-factor of odd rank.

Remark 3.7. Propositions 3.2 and 3.6 apply trivially to the case of odd n.

Example 3.1. We determine the S-admissible Howe subgroups of $\mathrm{O}(2)$ and $\mathrm{O}(4)$, see Figure 1. Cases (A) and (B) refer to Lemma 3.4.

$\mathrm{O}(2)$: O^1_2 belongs to case (A), $\mathrm{O}^1_1 \times \mathrm{O}^1_1$ belongs to case (B). Since O^1_2 is the centralizer of O^2_1, the only S-admissible Howe subgroup of $\mathrm{O}(2)$ is therefore U^1_1 which amounts to $\mathrm{SO}(2)$ itself. This is consistent with the fact that $\mathrm{SO}(2)$ is abelian.

$\mathrm{O}(4)$: $\mathrm{O}^1_2 \times \mathrm{O}^2_1$ and $\mathrm{O}^1_2 \times \mathrm{U}^1_1$ belong to case (A). $\mathrm{O}^1_1 \times \mathrm{O}^1_1 \times \mathrm{U}^1_1$, $\mathrm{O}^1_1 \times \mathrm{O}^1_1 \times \mathrm{O}^2_1$ and $\mathrm{O}^1_1 \times \mathrm{O}^1_1 \times \mathrm{O}^3_1$ belong to case (B). By cancelling these subgroups and their centralizers we arrive at the following list of S-admissible Howe subgroups of $\mathrm{O}(4)$: O^4_1, $\mathrm{O}^2_1 \times \mathrm{O}^2_1$, U^2_1, $\mathrm{O}^1_1 \times \mathrm{O}^1_1 \times \mathrm{O}^1_1 \times \mathrm{O}^1_1$, O^2_2, $\mathrm{U}^1_1 \times \mathrm{U}^1_1$, Sp^1_1, U^1_2, $\mathrm{O}^1_2 \times \mathrm{O}^1_2$, O^1_4.

It remains to discuss the passage to conjugacy classes in $\mathrm{SO}(n)$, n even. We will show

Proposition 3.8. *Let H be an* S*-admissible Howe subgroup H of* $\mathrm{O}(n)$*, n even. On intersection with* $\mathrm{SO}(n)$*, the* $\mathrm{O}(n)$*-conjugacy class of H passes to a single conjugacy class of Howe subgroups of* $\mathrm{SO}(n)$ *exactly in the following cases:*
– *H contains an* O*-factor of odd rank or odd multiplicity.*
– *H contains a* U*-factor of odd rank and odd multiplicity.*
Otherwise, the $\mathrm{O}(n)$*-conjugacy class of H gives rise to two distinct* $\mathrm{SO}(n)$*-conjugacy classes, generated by* $\mathrm{S}H$ *and* $\mathrm{S}(aHa^{-1})$ *for some $a \in$* $\mathrm{O}(n)$ *with negative determinant.*

We start with some preliminary observations. Since $\mathrm{SO}(n)$ is normal in $\mathrm{O}(n)$, intersection with $\mathrm{SO}(n)$ commutes with conjugation under either $\mathrm{O}(n)$ and $\mathrm{SO}(n)$. It is therefore sufficient to show that the conjugacy classes of H under $\mathrm{O}(n)$ and $\mathrm{SO}(n)$ coincide if and only if H belongs to one of the two cases listed in the proposition. If the two conjugacy classes do not coincide, it is clear that the $\mathrm{O}(n)$-class splits into two distinct $\mathrm{SO}(n)$-classes given by H and aHa^{-1}, where a is an arbitrary element of $\mathrm{O}(n)$ with $\det(a) = -1$. Since the conjugacy classes of H under $\mathrm{O}(n)$ and $\mathrm{SO}(n)$ coincide iff $\mathrm{N}_{\mathrm{O}(n)}(H)$ contains an element with negative determinant, we have to determine the normalizer of H. This will be done in two lemmas. We include the case of the identity connected component H_0 because this will be needed in Section 7 and we get it for granted here.

Lemma 3.9. *The normalizer of a Howe subgroup $H = \mathrm{I}_{\mathbb{K}_1\, m_i}^{\;k_i} \times \cdots \times \mathrm{I}_{\mathbb{K}_r\, m_r}^{\;k_r}$ of* $\mathrm{O}(n)$ *consists of all elements of* $\mathrm{O}(n)$ *which can be written in the form ab where*

$$a = \big(a_1 \oplus \overset{k_1}{\cdots} \oplus a_1\big) \oplus \cdots \oplus \big(a_r \oplus \overset{k_r}{\cdots} \oplus a_r\big)$$

with $a_i \in \mathrm{N}_{\mathrm{O}(\delta_i m_i)}\big(\mathrm{I}_{\mathbb{K}_i\, m_i}^{\;1}\big)$ and $b \in \mathrm{C}_{\mathrm{O}(n)}(H)$. The assertion still holds if H is replaced by H_0 and $\mathrm{I}_{\mathbb{K}_i\, m_i}^{\;k_i}$ by $\big(\mathrm{I}_{\mathbb{K}_i\, m_i}^{\;k_i}\big)_0$.

Proof. It is evident that any element of $\mathrm{O}(n)$ of the form given in the lemma belongs to the normalizer of H in $\mathrm{O}(n)$. Conversely, let $c \in \mathrm{N}_{\mathrm{O}(n)}(H)$. Consider the standard decomposition

$$\mathbb{R}^n = \big(\mathbb{R}^{\delta_1 m_1} \oplus \overset{k_1}{\cdots} \oplus \mathbb{R}^{\delta_1 m_1}\big) \oplus \cdots \oplus \big(\mathbb{R}^{\delta_r m_r} \oplus \overset{k_r}{\cdots} \oplus \mathbb{R}^{\delta_r m_r}\big)$$

into H-irreducible subspaces, used in the definition of H according to (6). The linear transformation c maps this decomposition to another orthogonal decomposition of $\mathbb{R}^n$ into H-irreducible subspaces. As abstract direct sums of orthogonal representations of H, the two decompositions are isometrically isomorphic. Hence, there exists $b_1 \in \mathrm{O}(n)$ commuting with H such that cb_1

leaves invariant each H-irreducible subspace $\mathbb{R}^{\delta_i m_i}$ in the standard decomposition above separately. Then

$$cb_1 = (c_{11} \oplus \cdots \oplus c_{1k_1}) \oplus \cdots \oplus (c_{r1} \oplus \cdots \oplus c_{rk_r})$$

with $c_{ij} \in \mathrm{N}_{\mathrm{O}(\delta_i m_i)}(\mathrm{I}^{\,1}_{\mathbb{K}_i\, m_i})$. Since the action of conjugation by c_{ij} on the operators of $\mathrm{I}^{\,1}_{\mathbb{K}_i\, m_i}$ is independent of j, $c_{ij} = a_i b_{ij}$, where $a_i \in \mathrm{N}_{\mathrm{O}(\delta_i m_i)}(\mathrm{I}^{\,1}_{\mathbb{K}_i\, m_i})$ and $b_{ij} \in \mathrm{C}_{\mathrm{O}(\delta_i m_i)}(\mathrm{I}^{\,1}_{\mathbb{K}_i\, m_i})$. Setting $b = (b_{11} \oplus \cdots \oplus b_{1k_1}) \oplus \cdots \oplus (b_{r1} \oplus \cdots \oplus b_{rk_r})\, b_1^{-1}$ we arrive at the assertion. The argument for H_0 is completely analogous. □

Lemma 3.10. *The normalizers* $\mathrm{N}_{\mathrm{O}(\delta_i m_i)}\big(\mathrm{I}^{\,1}_{\mathbb{K}_i\, m_i}\big)$ *are as follows.*
$\mathbb{K}_i = \mathbb{R}$*:* $\mathrm{N}_{\mathrm{O}(m_i)}\big(\mathrm{O}^1_{m_i}\big) = \mathrm{N}_{\mathrm{O}(m_i)}\big(\mathrm{SO}^1_{m_i}\big) = \mathrm{O}(m_i)\,.$
$\mathbb{K}_i = \mathbb{C}$*:* $\mathrm{N}_{\mathrm{O}(2m_i)}\big(\mathrm{U}^1_{m_i}\big)$ *is generated by* $\mathrm{U}^1_{m_i}$*, its centralizer and* $\mathbb{1}_{m_i,m_i} = \begin{bmatrix} \mathbb{1}_{m_i} & 0 \\ 0 & -\mathbb{1}_{m_i} \end{bmatrix}.$
$\mathbb{K}_i = \mathbb{H}$*:* $\mathrm{N}_{\mathrm{O}(4m_i)}\big(\mathrm{Sp}^1_{m_i}\big)$ *is generated by* $\mathrm{Sp}^1_{m_i}$ *and its centralizer.*
With the exception of the subgroups $\mathrm{O}^1_{m_i}$ *and* U^1_2*, in any case the induced homomorphism from the normalizer to the automorphism group of* $\mathrm{I}_{\mathbb{K}_i}(m_i)$ *or* $\mathrm{I}_{\mathbb{K}_i}(m_i)_0$ *is surjective. In the case of the subgroup* U^1_2*, the image of this homomorphism is generated by inner automorphisms and the outer automorphism of complex conjugation of matrices.*

Proof. For $\mathbb{K}_i = \mathbb{R}$, the normalizers are obvious. As for the automorphism groups it is known that for odd m_i all automorphisms of $\mathrm{O}(m_i)$ are inner whereas for even m_i they are generated by inner automorphisms and the outer automorphism $D : a \mapsto \det(a)a$. The automorphisms of $\mathrm{SO}(m_i)$ are restrictions of automorphisms of $\mathrm{O}(n)$. Since in case m_i is even, D acts trivially on $\mathrm{SO}(m_i)$, they are restrictions of inner automorphisms of $\mathrm{O}(n)$, as asserted.
For $\mathbb{K} = \mathbb{C}$, let $b \in \mathrm{N}_{\mathrm{O}(2m_i)}(\mathrm{U}^1_{m_i})$. Conjugation by b defines an automorphism of $\mathrm{U}(m_i)$. In case $m_i \neq 2$, the automorphism group of $\mathrm{U}(m_i)$ is generated by inner automorphisms and by the outer automorphism of complex conjugation of matrices. In case of $\mathrm{U}(2)$ there is one further generator, given by the outer automorphism $D(a) = \overline{\det a} \cdot a$. When $\mathrm{U}(m_i)$ is embedded into $\mathrm{O}(2m_i)$ via $\varphi_{\mathbb{C},\mathbb{R}}$, complex conjugation of matrices can be represented by conjugation by $\mathbb{1}_{m_i,m_i}$, but D can not. Thus, $b = Tb_1b_2$ where $T = \mathbb{1}_{2m_i}$ or $\mathbb{1}_{m_i,m_i}$, $b_1 \in \mathrm{U}^1_{m_i}$ and $b_2 \in \mathrm{C}_{\mathrm{O}(2m_i)}\big(\mathrm{U}^1_{m_i}\big)$. Conversely, any element of $\mathrm{O}(2m_i)$ of this form normalizes $\mathrm{U}^1_{m_i}$.
For $\mathbb{K}_i = \mathbb{H}$ the argument is similar. Here all automorphisms are inner. □

Proof of Proposition 3.8. According to the above considerations, we have to show that $\mathrm{N}_{\mathrm{O}(n)}(H)$ contains an element with negative determinant if and only if H belongs to one of the two cases given in the proposition. In the first case, H itself or its centralizer contains an element of negative determinant. In the second case, by Lemmas 3.9 and 3.10, $\mathrm{N}_{\mathrm{O}(n)}(H)$ contains an element which in the notation of Lemma 3.9 is given by ab with $b = \mathbb{1}_n$, $a_i = \mathbb{1}_{m_i,m_i}$ for the relevant U-factor and $a_i = \mathbb{1}_{\delta_i m_i}$ for the other factors. The determinant is $\det(ab) = (-1)^{m_i k_i} = -1$.
Conversely, let an element of $\mathrm{N}_{\mathrm{O}(n)}(H)$ with negative determinant be given. We can write it in the form ab where $b \in \mathrm{C}_{\mathrm{O}(n)}(H)$ and a is given in Lemma 3.9. Then

$$\det(ab) = (\det a_1)^{k_1} \cdots (\det a_r)^{k_r} \det b = -1\,.$$

If $\det b = -1$, $\mathrm{C}_{\mathrm{O}(n)}(H)$ contains an O-factor of odd multiplicity, hence H contains an O-factor of odd rank and we are in the first case of the proposition. If $\det b = 1$, one of the a_i must have negative determinant and the corresponding multiplicity k_i must be odd. By Lemma 3.10, if a_i has negative determinant then $\mathbb{K}_i = \mathbb{R}$, where we are in the first case of the proposition, or $\mathbb{K}_i = \mathbb{C}$ and m_i odd, where we are in the second case of the proposition. □



Figure 3: Hasse diagram of the set of conjugacy classes of Howe subgroups of SO(4). For the notation, see Figure 1.

To define a standard form for Howe subgroups of SO(n), we fix $a_{(n)} \in \mathrm{O}(n)$ with $\det(a_{(n)}) = -1$, e.g. $a_{(n)} = \begin{bmatrix} -1 & 0 \\ 0 & \mathbb{1}_{n-1} \end{bmatrix}$. For a Howe subgroup H of O(n) let

$$\mathrm{S}H^+ := \mathrm{S}H\,, \qquad \mathrm{S}H^- := a_{(n)}^{-1}\mathrm{S}H a_{(n)}\,. \tag{15}$$

By a subgroup of SO(n) of type [S$H^\pm$] we mean a subgroup of the form S$H^\pm$ where H is a Howe subgroup of O(n) in standard form. For simplicity, when treating SO(n) in the following, all results will be derived for both these types of subgroups, irrespective of that some of them may not be Howe or that SH^+ and SH^- may define the same conjugacy class in SO(n). If the latter is true we may just forget about SH^- and consider SH^+ alone. This applies in particular when n is odd.

Example 3.2. Consider the S-admissible Howe subgroups of O(2) and O(4) derived in Example 3.1. We determine their conjugacy classes in SO(2) and SO(4), respectively. For O(2), the situation is trivial. Let us check consistency, anyhow. The only S-admissible Howe subgroup of O(2) is U_1^1; it possesses a U-factor of odd rank and odd multiplicity and hence defines a single conjugacy class in SO(2) by Proposition 3.8 (consisting of SO(2) itself). For O(4), the S-admissible Howe subgroups which give rise to two distinct conjugacy classes in SO(4) by Proposition 3.8 are U_1^2, U_2^1, O_2^2 and Sp_1^1. Thus, there are altogether 14 conjugacy classes of Howe subgroups in SO(4). The corresponding Hasse diagram is shown in Figure 3. It is interesting to note that one of the two classes of type Sp_1^1 corresponds to the subgroup of left isoclinic rotations in SO(4) and the other one to the subgroup of right isoclinic rotations. These two subgroups are known to be conjugate in O(4) but not in SO(4), indeed.

Remark 3.11. O(n) contains Howe subgroups which give rise to two distinct conjugacy classes in SO(n) if and only if n is a multiple of 4. Indeed, any O-factor of such a Howe subgroup must have even rank and even multiplicity and any U-factor must have even rank or even multiplicity. Hence, any irreducible factor contributes to the dimension by a multiple of 4. Conversely, if $n = 4l$ then e.g. the Howe subgroup O_{2l}^2 gives rise to two distinct conjugacy classes in SO(n) by Proposition 3.8.

4 Principal bundles with a Howe subgroup as structure group

The aim of this section is to classify principal bundles whose structure group is a Howe subgroup of $\mathrm{O}(n)$, $\mathrm{SO}(n)$ or $\mathrm{Sp}(n)$ and whose base space is a simply connected CW-complex of dimension 4 without 2-torsion in $H^4(X,\mathbb{Z})$. We will use classifying spaces and classifying maps to construct characteristic classes which distinguish the bundles, see Appendices A and D for a brief overview. We start with introducing some notation. For a Lie group G, let $\mathrm{B}\,G$ denote the classifying space and let $[X,\mathrm{B}\,G]$ denote the set of homotopy classes of maps from X to $\mathrm{B}\,G$ or, equivalently, the set of isomorphism classes of principal G-bundles over X.
Let H be a Howe subgroup of $G=\mathrm{O}(n)$ or $G=\mathrm{Sp}(n)$. Due to (7) and (10), as a Lie group,

$$H \cong \mathrm{I}_{\mathbb{K}_1}(m_1) \times \cdots \times \mathrm{I}_{\mathbb{K}_r}(m_r)\,.$$

Accordingly, the classifying space of H decomposes as

$$\mathrm{B}\,H = \mathrm{B}\,\mathrm{I}_{\mathbb{K}_1}(m_1) \times \cdots \times \mathrm{B}\,\mathrm{I}_{\mathbb{K}_r}(m_r)$$

and any principal H-bundle Q over X decomposes as

$$Q = Q_1 \oplus \cdots \oplus Q_r$$

(fiber product), where the Q_i are $\mathrm{I}_{\mathbb{K}_i}(m_i)$-bundles over X. We refer to them as the factors of Q. More precisely, extending the terminology used for the Howe subgroups, we speak of O-, U- and Sp-factors, depending on whether $\mathbb{K}_i=\mathbb{R}$, $\mathbb{C}$ or $\mathbb{H}$, respectively. We will also have to consider the identity connected component H_0 of H. As a Lie group,

$$H_0 \cong \mathrm{I}_{\mathbb{K}_1}(m_1)_0 \times \cdots \times \mathrm{I}_{\mathbb{K}_r}(m_r)_0\,,$$

where $\mathrm{I}_{\mathbb{K}_i}(m_i)_0=\mathrm{SO}(m_i)$ for $\mathbb{K}_i=\mathbb{R}$ and $\mathrm{I}_{\mathbb{K}_i}(m_i)_0=\mathrm{I}_{\mathbb{K}_i}(m_i)$ for $\mathbb{K}_i=\mathbb{C}$, $\mathbb{H}$. The classifying space is

$$\mathrm{B}\,H_0 = \mathrm{B}\,\mathrm{I}_{\mathbb{K}_1}(m_1)_0 \times \cdots \times \mathrm{B}\,\mathrm{I}_{\mathbb{K}_r}(m_r)_0$$

and principal H_0-bundles over X decompose as

$$Q_0 = Q_{01} \times_X \cdots \times_X Q_{0r}\,,$$

where the Q_{0i} are $\mathrm{I}_{\mathbb{K}_i}(m_i)_0$-bundles over X, referred to as the factors of Q_0. Similarly to the Howe subbundles we speak of SO-, U- or Sp-factors here. Since X is simply connected, any H-bundle or $\mathrm{S}H^+$-bundle (in the case $G=\mathrm{O}(n)$) can be reduced to H_0, where the reductions are given by the connected components of the given H-bundle. As a consequence, the classification of H-bundles and $\mathrm{S}H^+$-bundles can be derived from the classification of H_0-bundles. To classify H_0-bundles it suffices to classify bundles with the factors of H_0 as structure group, i.e., $\mathrm{SO}(m)$, $\mathrm{U}(m)$ or $\mathrm{Sp}(m)$. The cases of $\mathrm{U}(m)$ and $\mathrm{Sp}(n)$ are standard; the necessary results will be cited below. Thus, the problem is reduced to the classification of principal $\mathrm{SO}(n)$-bundles. Since the solution is obvious for $n=1,2$, it remains to solve the problem for $n\geq 3$. We use methods introduced by Woodward [48] and improved by Čadek and Vanžura [8, 9, 10]. Since X has dimension 4, we have stability beginning with $n=5$ i.e. $[X,\mathrm{B}\,\mathrm{SO}(5)]=[X,\mathrm{B}\,\mathrm{SO}(6)]=\cdots=[X,\mathrm{B}\,\mathrm{SO}]$. Since X is simply connected, $[X,\mathrm{B}\,\mathrm{SO}]=[X,\mathrm{B}\,\mathrm{O}]$.

Remark 4.1. The reason for the assumption that X be simply connected is that it allows to reduce the study of $[X,\mathrm{B}\,\mathrm{O}(4)]$ to the study of $[X,\mathrm{B}\,\mathrm{SO}(4)]$. As we have to make it anyhow, we can use it further to make simplify arguments by, e.g., treating H-bundles and $\mathrm{S}H$-bundles simultaneously through H_0-bundles or switching freely between $\mathrm{O}(m)$-bundles and $\mathrm{SO}(m)$-bundles. The assumption of simple connectedness is however not as general as possible; in fact it is sufficient that $H^1(X,\mathbb{Z}_2)=0$.

4.1 Principal bundles with structure group $\mathrm{O}(n)$ or $\mathrm{SO}(n)$

Let $\theta_* : H^*(X,\mathbb{Z}_2) \to H^*(X,\mathbb{Z}_4)$ be the map induced by the inclusion homomorphism $\theta : \mathbb{Z}_2 \to \mathbb{Z}_4$ and let ρ and ρ_n denote the maps induced by reduction $\mathbb{Z}_4 \to \mathbb{Z}_2$ and $\mathbb{Z} \to \mathbb{Z}_n$ repectively. Let $\mathfrak{P}$ and Sq^1 denote the cohomology operations of Pontryagin square and Steenrod square, repectively, see Appendix C. Define a mapping

$$\mathfrak{R} : H^1(X,\mathbb{Z}_2) \oplus H^2(X,\mathbb{Z}_2) \oplus H^4(X,\mathbb{Z}_2) \oplus H^4(X,\mathbb{Z}) \to H^4(X,\mathbb{Z}_4)$$

by

$$\mathfrak{R}(u_1,u_2,u_4,v_4) := \rho_4 v_4 - \mathfrak{P}u_2 + \theta_*(u_1 \mathrm{Sq}^1 u_2 + u_4)\,,$$

for all spaces X. Let $w_k \in H^k(\mathrm{B\,O},\mathbb{Z}_2)$ be the k-th Stiefel-Whitney class, $p_1 \in H^4(\mathrm{B\,O},\mathbb{Z})$ the first Pontryagin class and denote $\alpha = (w_1,w_2,w_4,p_1)$. It can be shown, see Theorem D.3, that

$$\mathfrak{R}(\alpha) = 0 \tag{16}$$

holds for characteristic classes of real vector bundles of arbitrary dimension n. For any CW-complex X the 4-tuple α of cohomology elements defines a mapping

$$\alpha_* : [X,\mathrm{B\,O}] \to H^1(X,\mathbb{Z}_2) \oplus H^2(X,\mathbb{Z}_2) \oplus H^4(X,\mathbb{Z}_2) \oplus H^4(X,\mathbb{Z})$$

by

$$\alpha_*(\xi) := \xi^*(\alpha)\,, \quad \xi \in [X,\mathrm{B\,O}]\,. \tag{17}$$

Obviously, we have $\alpha_*(\xi) = \big(w_1(\xi),w_2(\xi),w_4(\xi),p_1(\xi)\big)$.

Theorem 4.2. *Let X be a CW-complex of dimension four. The mapping α_* defined by* (17) *is injective if $H^4(X,\mathbb{Z})$ has no element of order 4.*

Remark 4.3. *Let X be a 4-dimensional CW-complex without 4-torsion in $H^4(X,\mathbb{Z})$. Then, according to the above theorem the sets $[X,\mathrm{B\,O}(n)]$, $n \geq 5$, are classified by tuples (w_1,w_2,w_4,p_1). Moreover, via the natural projection $\mathrm{B\,SO}(n) \to \mathrm{B\,O}(n)$ we obtain a classification of $[X,\mathrm{B\,SO}(n)]$, $n \geq 5$, given by (w_2,w_4,p_1).*

Proof of Theorem 4.2: The strategy will be to replace $\mathrm{B\,O}$ by a 5-equivalent space. For that purpose, consider the following product of Eilenberg-MacLane spaces [2]

$$K = K(\mathbb{Z}_2,1) \times K(\mathbb{Z}_2,2) \times K(\mathbb{Z}_2,4) \times K(\mathbb{Z},4)\,.$$

By virtue of the natural isomorphism $H^n(\mathrm{B\,O},\cdot) = [\mathrm{B\,O},K(\cdot,n)]$, see Appendix B, the 4-tuple α of cohomology elements induces a mapping $\mathrm{B\,O} \to K$, which will be also denoted by the letter α.

Let

$$(\kappa_1,\kappa_2,\kappa_4,\chi_4) \in H^1(K(\mathbb{Z}_2,1),\mathbb{Z}_2) \oplus H^2(K(\mathbb{Z}_2,2),\mathbb{Z}_2) \oplus H^4(K(\mathbb{Z}_2,4),\mathbb{Z}_2) \oplus H^4(K(\mathbb{Z},4),\mathbb{Z})$$

denote the fundamental classes and let $(u_1,u_2,u_4,v_4) \in H^1(X,\mathbb{Z}_2) \oplus H^2(X,\mathbb{Z}_2) \oplus H^4(X,\mathbb{Z}_2) \oplus H^4(X,\mathbb{Z})$. Then via the isomorphism $H^n(X,\cdot) \to [X,K(\cdot,n)]$ we get a map $f = \varphi_1 \times \varphi_2 \times \varphi_4 \times \phi_4 \in [X,K]$ such that

$$(u_1,u_2,u_4,v_4) = f^*(\kappa_1 \otimes 1 \otimes 1 \otimes 1, 1 \otimes \kappa_2 \otimes 1 \otimes 1, 1 \otimes 1 \otimes \kappa_4 \otimes 1, 1 \otimes 1 \otimes 1 \otimes \chi_4).$$

---

[2] See Appendix B.

In what follows we leave out the 1's in the formulae. Since $\mathfrak{R}$ is built of cohomology operations we have:

$$\mathfrak{R} \circ f^*(\kappa_1, \kappa_2, \kappa_4, \chi_4) = f^* \circ \mathfrak{R}(\kappa_1, \kappa_2, \kappa_4, \chi_4).$$

That means $\mathfrak{R}$ is uniquely determined by its value on the fundamental classes. Moreover, since $\mathfrak{R}(\kappa_1, \kappa_2, \kappa_4, \chi_4) \in H^4(K, \mathbb{Z}_4)$, it defines a unique mapping $k : K \to K(\mathbb{Z}_4, 4)$ by

$$k^*(\iota_4) = \mathfrak{R}(\kappa_1, \kappa_2, \kappa_4, \chi_4)\,, \tag{18}$$

where $\iota_4 \in H^4(K(\mathbb{Z}_4, 4), \mathbb{Z}_4)$ denotes the fundamental class.

By passing to a homotopy equivalent K we can turn k into a fibration [19]. The homotopy fiber is given by $F = k^{-1}(*)$, where $*$ denotes the base point of $K(\mathbb{Z}_4, 4)$. We show that equation (16) implies that $k \circ \alpha$ is nullhomotopic:

$$\begin{aligned}
(k \circ \alpha)^*(\iota_4) &= \alpha^* \circ k^*(\iota_4) \\
&= \alpha^* \circ \mathfrak{R}(\kappa_1, \kappa_2, \kappa_4, \chi_4) \\
&= \mathfrak{R} \circ \alpha^*(\kappa_1, \kappa_2, \kappa_4, \chi_4) \\
&= \mathfrak{R}(\alpha^*\kappa_1, \alpha^*\kappa_2, \alpha^*\kappa_4, \alpha^*\chi_4) \\
&= \mathfrak{R}(\alpha) = 0\,,
\end{aligned}$$

where we have used again the isomorphism $H^n(X, \cdot) = [X, K(\cdot, n)]$. By lifting the homotopy $k \circ \alpha \sim 0$ to K we can achieve that $\alpha(\mathrm{B\,O}) \subseteq F$. Thus, there exists a lift $\tilde{\alpha} : \mathrm{B\,O} \to F$ and we have the following commutative diagram

$$\begin{tikzcd}
F \arrow[r, "j"] & K \arrow[r, "k"] & K(\mathbb{Z}_4, 4) \\
& \mathrm{B\,O} \arrow[u, "\alpha"] \arrow[ul, dashed, "\tilde{\alpha}"] &
\end{tikzcd} \tag{19}$$

Lemma 4.4. *The mapping $\tilde{\alpha}$ is a 5-equivalence.*

Proof of the Lemma: We will use that $j_* \circ \tilde{\alpha}_* = \alpha_*$ and determine $\tilde{\alpha}_*$ from j_* and α_*. The low-dimensional homotopy groups of $\mathrm{B\,O}$ are

$$\pi_1(\mathrm{B\,O}) = \pi_2(\mathrm{B\,O}) = \mathbb{Z}_2\,, \qquad \pi_3(\mathrm{B\,O}) = 0\,, \qquad \pi_4(\mathrm{B\,O}) = \mathbb{Z}\,.$$

The homotopy groups of K are

$$\pi_1(K) = \pi_2(K) = \mathbb{Z}_2\,, \qquad \pi_4(K) = \mathbb{Z}_2 \oplus \mathbb{Z}\,, \qquad \pi_i(K) = 0\,, \quad i = 3, i \geq 5\,.$$

The exact homotopy sequence of the fibration (19) decomposes into the portions

$$0 \to \pi_i(F) \xrightarrow{j_*} \pi_i(K) \xrightarrow{k_*} 0\,, \qquad i \neq 3, 4\,,$$

and

$$0 \to \pi_4(F) \xrightarrow{j_*} \pi_4(K) \xrightarrow{k_*} \pi_4(K(\mathbb{Z}_4, 4)) \to \pi_3(F) \xrightarrow{j_*} 0\,. \tag{20}$$

It follows that j_* is an isomorphism for all $i \neq 3, 4$ and injective for $i = 4$. First, this implies $\pi_5(F) = \pi_5(K) = 0$, hence $\tilde{\alpha}_*$ is onto for $i = 5$. Second, we conclude that for $i = 1, 2$, $\tilde{\alpha}_*$ is an isomorphism iff so is α_*.

We determine $\tilde{\alpha}_*$ in dimension $i = 3$ using homology. Let ν be an element of the subgroup $H_4(K(\mathbb{Z}, 4)) \subset H_4(K)$. Evaluation of (18) on $1 \times 1 \times 1 \times \nu$ yields for the l.h.s.

$$k^*(\iota_4)(1 \times 1 \times 1 \times \nu) = \iota_4(k_*(1 \times 1 \times 1 \times \nu))$$

and for the r.h.s.

$$\mathfrak{R}(\kappa_1, \kappa_2, \kappa_4, \chi_4)(1 \times 1 \times 1 \times \nu) = \rho_4 \circ \chi_4(\nu) .$$

Thus,

$$k_*(1 \times 1 \times 1 \times \nu) = {\iota_4}^{-1} \circ \rho_4 \circ \chi_4(\nu) .$$

Since χ_4 and ι_4 are isomorphisms and ρ_4 is surjective, k_* is surjective. Hence, $\pi_3(F) = 0$. Since also $\pi_3(\mathrm{B\,O}) = \pi_2(\mathrm{O}) = 0$, $\tilde{\alpha}_*$ is trivially an isomorphism in dimension $i = 3$.
Thus, it remains to show that α_* is an isomorphism in dimensions $i = 1, 2$ and ... in dimension $i = 4$. We have $\alpha_* = w_{1*} \times w_{2*} \times w_{4*} \times p_{1*}$. Thus, in dimension $i = 1, 2$ we get

$$\alpha_* = w_{i_*} : \pi_i(\mathrm{B\,O}) \to \pi_i(K(\mathbb{Z}_2, i))$$

and in dimension 4 we obtain

$$\alpha_* = w_{4*} \times p_{1*} : \pi_4(\mathrm{B\,O}) \to \pi_4(K(\mathbb{Z}_2, 4)) \times \pi_4(K(\mathbb{Z}, 4)) .$$

We calculate w_{i_*} for $i = 1, 2, 4$: Let $[\xi] \in \pi_i(\mathrm{B\,O}) = [S^i, \mathrm{B\,O}]$. Then $w_{i_*}([\xi]) = [w_i \circ \xi]$. By the isomorphism $[S^i, K(\mathbb{Z}_2, i)] \cong H^i(S^i, \mathbb{Z}_2)$ the element $[w_j \circ \xi]$ corresponds to

$$(w_i \circ \xi)^*(\kappa_i) = \xi^* \circ w_i^*(\kappa_i) = \xi^* w_i = w_i(\xi) ,$$

where $w_i(\xi)$ is the i-th Stiefel-Whitney class of the real vector bundle ξ . Since $\pi_i(K(\mathbb{Z}_2, i)) = \mathbb{Z}_2$ and since there exist real vector bundles with nonvanishing i-th Stiefel-Whitney class in all cases (the real, complex and quaternionic Hopf bundles), w_{i_*} is surjective. Since $\pi_1(\mathrm{B\,O}) = \mathbb{Z}_2 = \pi_2(\mathrm{B\,O})$ and $\pi_4(\mathrm{B\,O}) = \mathbb{Z}$ we conclude that w_{i_*} is an isomorphism for $i = 1, 2$ and w_{4_*} is reduction modulo 2 . This finishes the proof for $i = 1, 2$.
Next, we calculate p_{1*}: As above, for $[\xi] \in \pi_4(\mathrm{B\,O})$, we get

$$p_{1_*}([\xi]) = p_1(\xi) .$$

Using the standard relation $\rho_2 p_1 = w_2^2$, see Proposition D.4, together with $H^2(S^4, Z_2) = 0$, we conclude $\rho_2 p_1(\xi) = 0$. Thus, p_{1_*} is multiplication by an even number. Consider the quaternionic line bundle associated with the quaternionic Hopf bundle $S^3 \to S^7 \to S^4$, which can be viewed either as a real bundle ξ or as a complex bundle η . Then, $\xi^{\mathbb{C}} = \eta \oplus \eta$ and the third relation in Proposition D.5 together with the Whitney sum formula for the Chern classes yields

$$p_1(\xi) = -c_2(\xi^{\mathbb{C}}) = -2c_2(\eta) - c_1(\eta)^2 = -2c_2(\eta) .$$

Since $c_2(\eta)$ is a generator of $H^4(S^4, \mathbb{Z})$ we conclude that $(p_1)_*$ is multiplication by two. Hence, we get

$$\alpha_* = (w_4)_* \times (p_1)_* : \pi_4(\mathrm{B\,O}) = \mathbb{Z} \to \mathbb{Z}_2 \oplus \mathbb{Z} = \pi_4(K) , \quad n \mapsto (n \mod 2, 2n) .$$

In particular, α_* is injective. Since j_* is also injective and α_* and j_* have the same image, $\tilde{\alpha}_*$ is an isomorphism. □

We continue with the proof of the Theorem. Since $\tilde{\alpha}$ is a 5-equivalence the sets $[X, \mathrm{B\,O}]$ and $[X, F]$ are isomorphic. By iterating the process of forming homotopy fibers one obtains a sequence of fibrations associated with the fibration (19):

$$\ldots \longrightarrow \Omega F \longrightarrow \Omega K \longrightarrow \Omega B \longrightarrow F \longrightarrow K \longrightarrow B\,,$$

with $B = K(\mathbb{Z}_4, 4)$. Consider the following exact sequence related to this fibration (obtained by applying the functor $[X, -]$):

$$\cdots \longrightarrow [X, \Omega K] \xrightarrow{\Omega k_*} [X, K(\mathbb{Z}_4, 3)] \xrightarrow{i_*} [X, F] \xrightarrow{j_*} [X, K] \xrightarrow{k_*} [X, K(\mathbb{Z}_4, 4)]\,,$$

where $\Omega B = K(\mathbb{Z}_4, 3)$ has been used. The aim is to find a condition which ensures the injectivity of the map j_* in the sequence above. Obviously, j_* is injective if and only if i_* is the zero map. The latter is true if and only if Ωk_* is surjective. Thus, we have to calculate Ωk. We know that $\Omega k \in [\Omega K, K(\mathbb{Z}_4, 3)] = H^3(\Omega K, \mathbb{Z}_4)$. Under the isomorphism $H^3(K(\mathbb{Z}_4, 3), \mathbb{Z}_4) = [K(\mathbb{Z}_4, 3), K(\mathbb{Z}_4, 3)]$, the element $\iota_3 \in H^3(K(\mathbb{Z}_4, 3), \mathbb{Z}_4)$ obviously corresponds to the identity $[\mathrm{id}]$. Thus, $(\Omega k)^*(\iota_3)$ corresponds to $[\mathrm{id} \circ \Omega k] = [\Omega k]$ and computing Ωk is equivalent to computing $(\Omega k)^*(\iota_3)$. For this calculation consider the commutative diagram built from the path loop fibrations of K and $K(\mathbb{Z}_4, 4)$ respectively:

$$\begin{tikzcd}
\Omega K \arrow[r, "\Omega k"] \arrow[d] & K(\mathbb{Z}_4, 3) \arrow[d] \\
PK \arrow[r] \arrow[d] & PK(\mathbb{Z}_4, 4) \arrow[d] \\
K \arrow[r, "k"] & K(\mathbb{Z}_4, 4)
\end{tikzcd}$$

In what follows all suspension homomorphisms[3] occuring in the text will be denoted by σ. Using (76) together with $(\Omega k)^* \circ \sigma = \sigma \circ k^*$, see [30, Chapter 6.2], and (18) we get:

$$\begin{aligned}
(\Omega k)^*(\iota_3) &= (\Omega k)^*\big(\sigma(\iota_4)\big) \\
&= \sigma\big(k^*(\iota_4)\big) \\
&= \sigma\big(\mathfrak{R}(\kappa_1, \kappa_2, \kappa_4, \chi_4)\big) \\
&= \sigma\big(\rho_4 \chi_4 - \mathfrak{P}\kappa_2 + \theta_*(\kappa_1 \mathrm{Sq}^1 \kappa_2 + \kappa_4)\big)\,.
\end{aligned}$$

Obviously, Ωk_* is surjective if so is its restriction to $K(\mathbb{Z}, 3)$, which is the map induced by the first summand in the last line above. We have, see [30, Chapter 6.2],

$$\sigma \circ \rho_4(\chi_4) = \rho_4 \circ \sigma(\chi_4) = \rho_4(\chi_3)\,,$$

where $\chi_3 \in H^3(K(\mathbb{Z}, 3), \mathbb{Z})$ is the fundamental class and $K(\mathbb{Z}, 3) = \Omega K(\mathbb{Z}, 4) \subset \Omega K$. Now let $f \in [K(\mathbb{Z}, 3), K(\mathbb{Z}_4, 3)]$ denote the restriction of Ωk induced by $\rho_4(\chi_3)$, i.e.

$$f^*(\iota_3) = \rho_4(\chi_3)\,.$$

On the level of cohomology equivalence classes that means

$$[\iota_3 \circ \Omega k]_{\mathrm{co}} = [\rho_4 \circ \chi_3]_{\mathrm{co}}\,. \tag{21}$$

---

[3] see Appendix B

We claim that the following diagram is commutative:

$$\begin{tikzcd}
{[X, K(\mathbb{Z}, 3)]} \arrow[r, "f_*"] \arrow[d] & {[X, K(\mathbb{Z}_4, 3)]} \arrow[d] \\
H^3(X; \mathbb{Z}) \arrow[r, "\rho_4"] & H^3(X; \mathbb{Z}_4)
\end{tikzcd}$$

Here the unnamed arrows are the natural isomorphisms, see Appendix B. Let $g \in [X, K(\mathbb{Z}, 3)]$. Using (21) we get

$$\rho_4 g^*(\chi_3) = [\rho_4 \circ \chi_3 \circ g]_{\mathrm{co}} = [\iota_3 \circ f \circ g]_{\mathrm{co}} = \big(f \circ g\big)^*(\iota_3)\,.$$

Thus, the above diagram is commutative, indeed. That means j_* is injective if the map

$$H^3(X, \mathbb{Z}) \to H^3(X, \mathbb{Z}_4):\ x \mapsto \rho_4 x$$

is surjective. For the final step consider the following portion of the Bockstein sequence associated to $\mathbb{Z} \xrightarrow{4} \mathbb{Z} \xrightarrow{\rho_4} \mathbb{Z}_4$:

$$\begin{tikzcd}
H^3(X, \mathbb{Z}) \arrow[r, "\rho_4"] & H^3(X, \mathbb{Z}_4) \arrow[r, "\beta"] & H^4(X, \mathbb{Z}) \arrow[r, "4"] & H^4(X, \mathbb{Z})
\end{tikzcd}$$

Thus, ρ_4 is surjective iff $\beta = 0$, that means iff multiplication by 4 is injective, that means iff $H^4(X, \mathbb{Z})$ has no elements of order 4. This finishes the proof.

□

Corollary 4.5. *The image of the mapping α_* defined by* (17) *coincides with the set of* 4*-tuples*

$$\beta = (u_1, u_2, u_4, v_4) \in H^1(X, \mathbb{Z}_2) \oplus H^2(X, \mathbb{Z}_2) \oplus H^4(X, \mathbb{Z}_2) \oplus H^4(X, \mathbb{Z})$$

fulfilling the relation

$$\mathfrak{R}(u_1, u_2, u_4, v_4) = 0\,. \tag{22}$$

Proof: Let β be a 4-tuple fulfilling relation (22) and let us denote the corresponding mappping $X \to K$ also by β. By the same arguments as above, there exists a lift $\tilde{\beta}$ of β, i.e. the following diagram commutes:

$$\begin{tikzcd}
F \arrow[r, "j"] & K \arrow[r, "k"] & K(\mathbb{Z}_4, 4) \\
 & X \arrow[u, "\beta"'] \arrow[ul, dashed, "\tilde{\beta}"] &
\end{tikzcd} \tag{23}$$

Putting together the diagrams (19) and (23) we obtain

$$\begin{tikzcd}
 & \mathrm{B\,O} \arrow[dl, "\tilde{\alpha}"'] \arrow[d, "\alpha"] & \\
F \arrow[r, "j"] & K \arrow[r, "k"] & K(\mathbb{Z}_4, 4) \\
 & X \arrow[u, "\beta"'] \arrow[ul, "\tilde{\beta}"] &
\end{tikzcd} \tag{24}$$

By Lemma 4.4, $\tilde{\alpha}$ induces an isomorphism

$$\tilde{\alpha}_*\colon [X,F] \to [X,\mathrm{B\,O}]\,.$$

Thus, using the inverse isomorphism $(\tilde{\alpha}_*)^{-1}$ we get a mapping class $\hat{\beta} := (\tilde{\alpha}_*)^{-1}(\tilde{\beta}) \in [X,\mathrm{B\,O}]$. We compute

$$\alpha \circ \hat{\beta} = j \circ \tilde{\alpha} \circ \hat{\beta} = j \circ \tilde{\alpha}_*\big((\tilde{\alpha}_*)^{-1}(\tilde{\beta})\big) = j \circ \tilde{\beta} = \beta\,,$$

or on the level of cohomology classes $\alpha_*(\hat{\beta}) = \beta$. □

Next, in analogy to the situation in Theorem 4.2 we consider the 3-tuple $\alpha' = (w_1, w_2, p_1)$ of cohomology elements with $w_k \in H^k(\mathrm{B\,O}(3),\mathbb{Z}_2)$ and $p_1 \in H^4(\mathrm{B\,O}(3),\mathbb{Z})$ being the Stiefel-Whitney classes and the Pontryagin class respectively. As above, α' defines the mapping

$$\alpha'_* : [X,\mathrm{B\,O}(3)] \to H^1(X,\mathbb{Z}_2) \oplus H^2(X,\mathbb{Z}_2) \oplus H^4(X,\mathbb{Z})\,, \quad \alpha'_*(\xi) := \xi^*(\alpha')\,, \tag{25}$$

for any CW-complex X. This means $\alpha'_*(\xi) = \big(w_1(\xi), w_2(\xi), p_1(\xi)\big)$.

Theorem 4.6. *Let X be a complex of dimension four. Then, the mapping α'_* defined by* (25) *is injective if $H^4(X,\mathbb{Z})$ has no element of order* 4.

Remark 4.7. *Let X be a* 4*-dimensional CW-complex without* 4*-torsion in $H^4(X,\mathbb{Z})$. Then, according to the above theorem the sets $[X,\mathrm{B\,O}(3)]$ are classified by tuples (w_1,w_2,p_1). Moreover, via the natural projection $\mathrm{B\,SO}(3) \to \mathrm{B\,O}(3)$ we gain a classification of $[X,\mathrm{B\,SO}(3)]$ given by (w_2,p_1).*

The proof of Theorem 4.6 is completely analogous to the proof of Theorem 4.2 and will be omitted. We also have the analogous

Corollary 4.8. *The image of the mapping α'_* defined by* (25) *coincides with the set of* 3*-tuples*

$$(u_1,u_2,v_4) \in H^1(X,\mathbb{Z}_2) \oplus H^2(X,\mathbb{Z}_2) \oplus H^4(X,\mathbb{Z})$$

fulfilling the relation

$$\rho_4 v_4 - \mathfrak{P} u_2 - \theta_*(u_1 \mathrm{Sq}^1 u_2) = 0\,. \tag{26}$$

The next result generalizes a classical one (see [45]).

Theorem 4.9. *Let X be a complex of even dimension n. Then the number of different isomorphism classes of n-dimensional orientable vector bundles which have the same Euler class and which are stably equivalent is bounded from above by the number of* 2*-torsion elements in $H^n(X,\mathbb{Z})$.*

Corollary 4.10. *If $H^n(X,\mathbb{Z})$ has no* 2*-torsion then two vector bundles ξ and η over X are isomorphic iff they are stably equivalent and have the same Euler class.*

Remark 4.11. For the case $n = 4$ and under the assumption that $H^4(X,\mathbb{Z})$ has no 2-torsion, it follows from Theorem 4.2 that two principal SO(4)-bundles are stably equivalent iff their second and fourth Stiefel-Whitney classes and their first Pontryagin classes coincide. Thus, principal SO(4)-bundles P over X are classified by tuples $(w_2(P), w_4(P), p_1(P), e(P))$. This follows from the fact that $(w_2(P), w_4(P), p_1(P)) = f_P^* \circ \mu^*(w_2, w_4, p_1)$, where $\mu : \mathrm{B\,SO}(4) \to \mathrm{B\,SO}$ is the natural projection, f_P is the classifying map of P and (w_2, w_4, p_1) are the Stiefel-Whitney classes and the first Pontryagin class in the cohomology of $\mathrm{B\,SO}$.

Proof of Theorem 4.9: We formulate the statement in terms of a lifting property: Let there be given 2 elements $f_i \in [X, \mathrm{B\,SO}(n)]$, $i = 1, 2$, which define stably equivalent bundles. Stable equivalence means $\pi \circ f_1 = \pi \circ f_2$, with π being the projection of the canonical fibration $S^n \hookrightarrow \mathrm{B\,SO}(n) \to \mathrm{B\,SO}(n+1)$. Thus, the question is how many lifts (defining the same Euler class) of a given map $f : X \to \mathrm{B\,SO}(n+1)$ do exist. This problem will be solved by using the Moore-Postnikov tower for the fibration $\mathrm{B\,SO}(n) \to \mathrm{B\,SO}(n+1)$ (see [44] for the details of this construction). It turns out that it is sufficient to consider its first nontrivial stage:

$$\begin{tikzcd}
 & S^n \arrow[d, "\iota'"] & K(\mathbb{Z}, n) \arrow[d, "\iota"] & \\
 & \mathrm{B\,SO}(n) \arrow[r, "\varphi"] \arrow[d, "\pi"] & E \arrow[d, "p"] & \\
X \arrow[r] \arrow[ur, shift left, "f_1"] \arrow[ur, shift right, "f_2"'] & \mathrm{B\,SO}(n+1) \arrow[r, equal] & \mathrm{B\,SO}(n+1) \arrow[r, "k"] & K(\mathbb{Z}, n+1)
\end{tikzcd}$$

For the convenience of the reader we recall that this diagram is commutative and has the following properties:

1. The map φ is an $(n+1)$-equivalence.
2. The fibration $p : E \to \mathrm{B\,SO}(n+1)$ is the pull-back of the path loop fibration of $K(\mathbb{Z}, n+1)$ under the mapping k .
3. The map $k : \mathrm{B\,SO}(n+1) \to K(\mathbb{Z}, n+1)$ is given by

$$k^*(\iota_{n+1}) = \tau'(o_n)\,, \tag{27}$$

where τ' is the transgression homomorphisms, see Appendix B, associated with the fibration $\pi : \mathrm{B\,SO}(n) \to \mathrm{B\,SO}(n+1)$, ι_{n+1} is the fundamental class of $K(\mathbb{Z}, n+1)$ and $o_n \in H^n(S^n, \mathbb{Z})$ is the n-th cohomology generator.

Lemma 4.12. *Let w_n be the n-th Stiefel-Whitney class of* $\mathrm{B\,SO}(n+1)$. *We have*

$$\tau'(o_n) = \beta(w_n)\,, \tag{28}$$

where β is the Bockstein homomorphism associated to $0 \to \mathbb{Z} \to \mathbb{Z} \to \mathbb{Z}_2 \to 0$.

Proof of the Lemma: Consider the following portion of the Serre exact sequence associated to the fibration $\pi : \mathrm{B\,SO}(n) \to \mathrm{B\,SO}(n+1)$:

$$\cdots \longrightarrow H^n(S^n, \mathbb{Z}) \xrightarrow{\tau'} H^{n+1}(\mathrm{B\,SO}(n+1), \mathbb{Z}) \xrightarrow{\pi^*} H^{n+1}(\mathrm{B\,SO}(n), \mathbb{Z}).$$

We have $\mathrm{im}\,(\tau') = \ker(\pi^*)$. Recall that n is even. According to (88) in Appendix D, the Euler class of $\mathrm{B\,SO}(n+1)$ satisfies $\rho_2 e = w_{n+1}$. On the other hand, we have

$$\rho_2 \circ \beta(w_n) = \mathrm{Sq}^1(w_n) = w_{n+1}\,.$$

Since ρ_2 is injective on the torsion elements of $H^*(\mathrm{B\,SO}(n+1), \mathbb{Z})$, we conclude that $e = \beta(w_n)$. Now consider the Gysin sequence of the fibration $S^n \hookrightarrow \mathrm{B\,SO}(n) \xrightarrow{\pi} \mathrm{B\,SO}(n+1)$:

$$\cdots \longrightarrow H^0(\mathrm{B\,SO}(n+1), \mathbb{Z}) \xrightarrow{\beta(w_n)} H^{n+1}(\mathrm{B\,SO}(n+1), \mathbb{Z}) \xrightarrow{\pi^*} H^{n+1}(\mathrm{B\,SO}(n), \mathbb{Z}) \longrightarrow \cdots .$$

Obviously, $\ker \pi^*$ is generated by $\beta(w_n)$ which proves the lemma. □

By point 1 of the above list, the induced mapping

$$\varphi_* : [X, \mathrm{B}\,\mathrm{SO}(n)] \to [X, E]$$

is an isomorhism. Thus, instead of f_i we consider their images under this isomorphism, which we denote by the same letter. By point 2 as well as the above Lemma we have

$$E = \{(b, \gamma) \in \mathrm{B}\,\mathrm{SO}(n+1) \times PK(\mathbb{Z}, n+1) \mid \beta(w_n)(b) = \gamma(1)\}\,.$$

Since $K(\mathbb{Z}, n) = \Omega K(\mathbb{Z}, n+1)$ we can define a multiplication $m : E \times K(\mathbb{Z}, n) \to E$ by

$$m\big((b, \gamma), \alpha\big) = (b, \gamma \circ \alpha)\,.$$

In every point $x \in X$ the two lifts f_1 and f_2 differ only by an element $\alpha \in \Omega K(\mathbb{Z}, n{+}1) = K(\mathbb{Z}, n)$. Thus, there is a map $d : X \to K(\mathbb{Z}, n)$ such that

$$f_2 = m(f_1, d)\,, \tag{29}$$

where m is viewed as point-wise product.

Lemma 4.13. *The induced mapping* $m^* : H^n(E, \mathbb{Z}) \to H^n(E \times \Omega K(\mathbb{Z}, n+1))$ *is given by*

$$m^*(x_n) = x_n \otimes 1 + 1 \otimes \iota^*(x_n)\,, \tag{30}$$

where ι *is the inclusion of the fiber into the fibration* $\Omega K(\mathbb{Z}, n+1) \hookrightarrow E \to \mathrm{B}\,\mathrm{SO}(n+1)$*.*

Proof of the Lemma: Using the Künneth Theorem we obtain

$$\begin{aligned} H^n(E \times \Omega K(\mathbb{Z}, n+1), \mathbb{Z}) = H^n(E, \mathbb{Z}) &\otimes H^0(\Omega K(\mathbb{Z}, n+1), \mathbb{Z}) \\ \oplus\, H^0(E, \mathbb{Z}) &\otimes H^n(\Omega K(\mathbb{Z}, n+1), \mathbb{Z}). \end{aligned}$$

Since $H^0(E, \mathbb{Z}) = \mathbb{Z} = H^0(\Omega K(\mathbb{Z}, n+1), \mathbb{Z})$ we get

$$\begin{aligned} m^*(x_n) &= (\sum_k (y_n)_k \otimes \mu_k) + (\sum_r \nu_r \otimes (z_n)_r) \\ &= (\sum_k \mu_k (y_n)_k \otimes 1) + (1 \otimes \sum_r \nu_r (z_n)_r) \\ &= (y_n \otimes 1) + (1 \otimes z_n)\,. \end{aligned}$$

For the computation of y_n and z_n let the canonical inclusions and projections of $E \times \Omega K(\mathbb{Z}, n{+}1)$ to its factors be denoted according to the following diagram:

$$\begin{tikzcd}
& E \times \Omega K(\mathbb{Z}, n+1) \arrow[dl, shift left, "pr_1"] \arrow[dr, shift left, "pr_2"] & \\
E \arrow[ur, shift left, "i_1"] & & \Omega K(\mathbb{Z}, n+1) \arrow[ul, shift left, "i_2"]
\end{tikzcd}$$

Obviously, we have $m \circ i_1 = \mathrm{id}$ and hence:

$$x_n = i_1^*\, m^*(x_n) = i_1^*(y_n \otimes 1 + 1 \otimes z_n) = y_n$$

where the last equality simply follows from the fact that $i_1^*\, pr_1^*(y_n) = y_n$ and $i_1^*\, pr_2^*(y_n) = 0$. We show that $z_n = \iota^*(x_n)$. For that purpose note that $m \circ i_2 = \iota$ and hence:

$$\iota^*(x_n) = i_2^*\, m^*(x_n) = i_2^*(x_n \otimes 1 + 1 \otimes z_n) = z_n\,.$$

□

>From Proposition E.1 applied to the map $\varphi : \mathrm{B}\,\mathrm{SO}(n) \to E$ we know that

$$\varphi^* : H^n(E,\mathbb{Z}) \to H^n(\mathrm{B}\,\mathrm{SO}(n),\mathbb{Z})$$

is an isomorphism. In what follows we identify the Euler class $e \in H^n(\mathrm{B}\,\mathrm{SO}(n),\mathbb{Z})$ with its image $(\varphi^*)^{-1}(e)$ in $H^n(E,\mathbb{Z})$.

Lemma 4.14. *Let* $\iota_n \in H^n(K(\mathbb{Z},n),\mathbb{Z})$ *be the fundamental class. Then we have*

$$\iota^*(e) = (\pm)2\iota_n\,.$$

Proof of the Lemma: Let us consider the following portion of the Serre sequence for the fibration $K(\mathbb{Z},n) \overset{\iota}{\hookrightarrow} E \overset{p}{\to} \mathrm{B}\,\mathrm{SO}(n+1)$ as well as for the fibration $S^n \overset{\iota'}{\hookrightarrow} \mathrm{B}\,\mathrm{SO}(n) \overset{\pi}{\to} \mathrm{B}\,\mathrm{SO}(n+1)$:

$$\begin{tikzcd}
H^n(\mathrm{B}\,\mathrm{SO}(n+1),\mathbb{Z}) \arrow[r, "p^*"] \arrow[d, equal] & H^n(E,\mathbb{Z}) \arrow[r, "\iota^*"] \arrow[d, "\varphi^*"] & H^n(K(\mathbb{Z},n),\mathbb{Z}) \arrow[d, "(\varphi\restriction_{S^n})^*"] \\
H^n(\mathrm{B}\,\mathrm{SO}(n+1),\mathbb{Z}) \arrow[r, "\pi^*"] & H^n(\mathrm{B}\,\mathrm{SO}(n),\mathbb{Z}) \arrow[r, "(\iota')^*"] & H^n(S^n,\mathbb{Z})
\end{tikzcd}$$

$$\begin{tikzcd}
H^n(K(\mathbb{Z},n),\mathbb{Z}) \arrow[r, "\tau"] \arrow[d, "(\varphi\restriction_{S_n})^*"] & H^{n+1}(\mathrm{B}\,\mathrm{SO}(n+1),\mathbb{Z}) \arrow[r, "p^*"] \arrow[d, equal] & H^{n+1}(E,\mathbb{Z}) \arrow[d, "\varphi^*"] \\
H^n(S^n,\mathbb{Z}) \arrow[r, "\tau'"] & H^{n+1}(\mathrm{B}\,\mathrm{SO}(n+1),\mathbb{Z}) \arrow[r, "\pi^*"] & H^{n+1}(\mathrm{B}\,\mathrm{SO}(n),\mathbb{Z})
\end{tikzcd}$$

Since $p\circ\varphi = \pi$ we have $\pi^* = \varphi^*\circ p^*$ and hence $\ker\pi^* = \ker\varphi^*\circ p^*$. Moreover, applying Proposition E.1 again to the mapping $\varphi : \mathrm{B}\,\mathrm{SO}(n) \to E$ we see that $\varphi^* : H^{n+1}(E,\mathbb{Z}) \to H^{n+1}(\mathrm{B}\,\mathrm{SO}(n),\mathbb{Z})$ is injective and hence $\ker\pi^* = \ker p^*$. Since the Serre sequence is exact this implies $\mathrm{im}\,\tau' = \mathrm{im}\,\tau$. Since both, $H^n(S^n,\mathbb{Z})$ and $H^n(K(\mathbb{Z},n),\mathbb{Z})$ are generated by a single element, namely o_n and ι_n, Lemma 4.12 implies

$$\tau'(o_n) = \beta(w_n) = \tau(\iota_n)\,.$$

Since $\beta(w_n)$ is a 2-torsion element the kernel of τ is given by the multiplication map by 2. Since the Sequence is exact this kernel equals $\mathrm{im}\,(\iota^*)$, i.e. the image of ι^* is generated by $2\iota_n$. It is well-known that $H^n(\mathrm{B}\,\mathrm{SO}(n),\mathbb{Z})/\mathrm{im}\,\pi^*$ is generated by the Euler class e. Since $\pi^* = \varphi^*\circ p^*$ we see that $H^n(E,\mathbb{Z})/\mathrm{im}\,p^*$ is generated by e, too. (Recall that we have identified e with its image under $(\varphi^*)^{-1}$.) Thus, we obtain the result $\iota^*(e) = \pm 2\iota_n$. □

For the final step of the proof we write down (29) more precisely as the composition:

$$\begin{tikzcd}
X \arrow[r, "\Delta"] \arrow[rrr, bend right, "f_2"'] & X \times X \arrow[r, "f_1\times d"] & E \times K(\mathbb{Z},n) \arrow[r, "m"] & E
\end{tikzcd}$$

and calculate:

$$\begin{aligned}
f_2^*(e) &= \Delta^*(f_1\times d)^* m^*(e) \\
&= \Delta^*(f_1\times d)^*\big(e\otimes 1 + 1\otimes\iota^*(e)\big) \\
&= \Delta^*(f_1\times d)^*(e\otimes 1 \pm 1\otimes 2\iota_n) \\
&= \Delta^*\big(f_1^*(e)\otimes 1 \pm 1\otimes 2d^*(\iota_n)\big) \\
&= f_1^*(e) \pm 2d^*(\iota_n)\,.
\end{aligned}$$

Assume that the two stably equivalent bundles f_1 and f_2 have the same Euler class. Then, the last equation implies that they can only differ by a map d with the property $2d = 0$. Realizing that the homotopy classes of such maps are in one-to-one correspondence with 2-torsion elements of $H^n(X,\mathbb{Z})$ it completes the proof of the theorem. □

Remark 4.15. The Stiefel Whitney class w_3 does not enter in any of the above classification results. This suggests that there should be a relation expressing w_3 in terms of the other characteristic classes. Indeed, (89) implies

$$w_3(P) = w_1 w_2 + \mathrm{Sq}^1 w_2(P)\,. \tag{31}$$

In particular, for X being simply connected or for principal $SO(n)$-bundles P we have $w_1(P) = 0$ and, therefore,

$$w_3(P) = \mathrm{Sq}^1 w_2(P)\,. \tag{32}$$

For further use, we will now formulate the classification of $\mathrm{SO}(n)$- and $\mathrm{O}(n)$-bundles over 4-dimensional CW-complexes X in a universal way. For convenience, in both cases we assume that X is simply connected, although this assumption is necessary only for the $\mathrm{O}(n)$-bundles[4].

Theorem 4.16. *Let X be a simply connected CW-complex of dimension 4 without 2-torsion in $H^4(X,\mathbb{Z})$. $\mathrm{SO}(n)$-bundles over X are classified by the characteristic classes w, p, and e. More precisely, the map*

$$\begin{aligned} [X, \mathrm{B}\,\mathrm{SO}(n)] &\to H^2(X,\mathbb{Z}_2)\times H^4(X,\mathbb{Z}_2)\times H^4(X,\mathbb{Z})\times H^n(X,\mathbb{Z}) \\ P &\mapsto \big(w_2(P), w_4(P), p_1(P), e(P)\big) \end{aligned} \tag{33}$$

is a bijection onto the subset defined by the fundamental relation

$$\mathfrak{R}(0, w_2(P), w_4(P), p_1(P)) \equiv \rho_4\, p_1(P) - \mathfrak{P} w_2(P) + \theta_* w_4(P) = 0 \tag{34}$$

and the following special relations:

$n = 1$: $w_2(P) = w_4(P) = p_1(P) = e(P) = 0$.

$n = 2$: $w_2(P) = \rho_2 e(P)$, $w_4(P) = 0$, $p_1(P) = e(P)^2$.

$n = 3$: $w_4(P) = 0$, $\rho_2 e(P) = \mathrm{Sq}^1 w_2(P)$.

$n = 4$: $w_4(P) = \rho_2 e(P)$.

Remark 4.17. In case $n = 3$, the relation for e determines $e(P)$ uniquely because $e(P)$ is a 2-torsion element here, see [20, Prop. 3.13]. Hence, the Euler class is a relevant parameter for $n = 2$ and $n = 4$ only.

Proof. We have $w_1(P) = 0$ and $w_3(P) = \mathrm{Sq}^1 w_2(P)$, by Remark 4.15. For the stable case $n \geq 5$, the theorem follows from Theorem 4.2 and Corollary 4.5. As mentioned earlier, see (16), the fundamental relation is fulfilled in all cases. For $n = 1$, the assertion is obvious. For $n = 2$ we have the isomorphism $\varphi_{\mathbb{C},\mathbb{R}} : \mathrm{U}(1) \to \mathrm{SO}(2)$. According to Proposition D.5, $\big(\mathrm{B}\,\varphi_{\mathbb{C},\mathbb{R}}\big)^* e = c_1$. Since c_1 defines a bijection from $[X, \mathrm{B}\,\mathrm{U}(1)]$ onto $H^2(X,\mathbb{Z})$, the same holds for e and $[X, \mathrm{B}\,\mathrm{SO}(2)]$. It remains to check the relations. The first one follows from the relation $\rho_2 e = w_2$, see (88). The second one is obvious. The third one follows from Theorem D.2.

---

[4] See however Remark 4.1

For $n = 3$, $w_4 = 0$. The relation for e is due to (31) and (88). Hence, the assertion follows from Theorem 4.6 and Corollary 4.8.

For $n = 4$, the classification is given by tuples $(w_2(P), w_4(P), p_1(P), e(P))$ of characteristic classes according to Theorem 4.2 and Corollary 4.10. It remains to show that the image of the mapping (33) is defined by the above relations: For that purpose, observe that for every SO(4)-bundle with classifying map f there exists a stable bundle $\mu \circ f$ induced via the natural projection $\mu : \mathrm{B\,SO}(4) \to \mathrm{B\,SO}$. For a given tuple (w_2, w_4, p_1), fulfilling the fundamental relation, there is always a stable bundle and hence an SO(4)-bundle P with classifying map f_P and classes $(w_2(P), w_4(P), p_1(P)) = f_P^* \circ \mu^*(w_2, w_4, p_1)$, which also fulfil the fundamental relation. This follows from Corollary 4.5, see also Remark 4.11. For any class $e \in H^4(X, \mathbb{Z})$ fulfilling the relation $w_4(P) = \rho_2 e$ we take $c_2 = \mathrm{B}\,\varphi_{\mathbb{C},\mathbb{R}}(e)$. This is the second Chern class of a principal U(2)-bundle with classifying map g. Then, $\mathrm{B}\,\varphi_{\mathbb{C},\mathbb{R}} \circ g$ defines a principal SO(4)-bundle with Euler class e. The relation $w_4(P) = \rho_2 e$ follows from (88). □

Since X is simply connected, from the classification of $[X, \mathrm{B\,SO}(n)]$ we obtain a classification of $[X, \mathrm{B\,O}(n)]$ by identifying SO(n)-bundles which are reductions of the same O(n)-bundle, i.e., which differ at most in the sign of the Euler class. To formalize that, for $\alpha \in H^k(X, \mathbb{Z})$ let $[\alpha]$ denote the class of α under the equivalence relation $\alpha \sim \beta \Leftrightarrow \alpha + \beta = 0$ and let $\mathrm{P}H^k(X, \mathbb{Z})$ denote the set of equivalence classes. On $\mathrm{P}H^*(X, \mathbb{Z})$, the following operations can be defined:

- the cup product by $[\alpha][\beta] = [\alpha\beta]$,
- the square as a map $\mathrm{P}H^k(X, \mathbb{Z}) \to H^{2k}(X, \mathbb{Z})$ by $[\alpha]^2 = \alpha^2$,
- reduction mod 2 as a map $\mathrm{P}H^*(X, \mathbb{Z}) \to H^*(X, \mathbb{Z}_2)$ by $\rho_2([\alpha]) = \rho_2\alpha$.

Define a map $[e] : [X, \mathrm{B\,O}(n)] \to \mathrm{P}H^n(X, \mathbb{Z})$ by $[e](P) := [e(P_0)]$, where P_0 is some reduction of P to SO(n). We will loosely speak of $[e]$ as a characteristic class for O(n)-bundles, although it is not a cohomology element of X and it is not defined by a cohomology element of $\mathrm{B\,O}(n)$. Passing to equivalence classes of $e(P)$ in Theorem 4.16 we obtain

Corollary 4.18. *Theorem 4.16 remains true when* SO(n) *is replaced by* O(n), $H^n(X, \mathbb{Z})$ *by* $\mathrm{P}H^n(X, \mathbb{Z})$ *and* $e(P)$ *by* $[e](P)$. □

4.2 Principal bundles with structure groups H_0, H and $\mathrm{S}H^{\pm}$

We recall the classification of U(n) and Sp(n)-bundles over a CW-complex X of dimension 4:

- The Chern class c of $\mathrm{B\,U}(n)$ defines a bijection from $[X, \mathrm{B\,U}(n)]$ onto $H^2(X, \mathbb{Z})$ for $n = 1$ and onto $H^2(X, \mathbb{Z}) \times H^4(X, \mathbb{Z})$ for $n \geq 2$.
- The Chern class c of $\mathrm{B\,Sp}(n)$, defined by $c\big(\mathrm{Sp}(n)\big) = \big(\mathrm{B}\,\varphi_{\mathbb{H},\mathbb{C}}\big)^* c\big(\mathrm{U}(2n)\big)$, defines a bijection from $[X, \mathrm{B\,Sp}(n)]$ onto $H^4(X, \mathbb{Z})$. This is a consequence of the isomorphy of Sp(1) and SU(2) and the fact that in 4 dimensions, Sp(1) is already the stable case.

Putting all of the above facts together we obtain the following classification result for Lie groups, which appear as Howe subgroups of O(n) and Sp(n).

Theorem 4.19. *Let X be a simply connected CW-complex of dimension 4 without 2-torsion in $H^4(X, \mathbb{Z})$ and let $H = \mathrm{I}_{\mathbb{K}_1}(m_1) \times \cdots \times \mathrm{I}_{\mathbb{K}_r}(m_r)$, where $\mathbb{K}_i = \mathbb{R}, \mathbb{C}, \mathbb{H}$. Principal bundles Q over X with structure group H_0 or H are classified by the collection of characteristic classes of their factors Q_i, i.e.,*

- *in case of H_0 by $w(Q_i)$, $p(Q_i)$ and $e(Q_i)$ for $\mathbb{K}_i = \mathbb{R}$ and $c(Q_i)$ for $\mathbb{K}_i = \mathbb{C}, \mathbb{H}$,*
- *in case of H by $w(Q_i)$, $p(Q_i)$ and $[e](Q_i)$ for $\mathbb{K}_i = \mathbb{R}$ and $c(Q_i)$ for $\mathbb{K}_i = \mathbb{C}, \mathbb{H}$.*

For $\mathbb{K}_i = \mathbb{R}$, the characteristic classes are subject to the relations given in Theorem 4.16*, Corollary* 4.18 *and Formula* (31). ☐

More precisely, the statement of the theorem reads as follows. Let $K = H_0$ or H and let J_i denote the image of the characteristic classes of the factor Q_i in the cohomology of X (where it is understood that in case $K = H$ the image of the Euler class is a subset of $\mathrm{P}H^{m_i}(M, \mathbb{Z})$). Then the collection of characteristic classes of the factors Q_i of Q defines a bijection from $[X, \mathrm{B}\, K]$ onto $J_1 \times \cdots \times J_r$.

It remains to treat the case of principal bundles whose structure group is a subgroup of type $[\mathrm{S}H^{\pm}]$ of $\mathrm{SO}(n)$. In the following we assume that H has an O-factor (otherwise $H \subseteq \mathrm{SO}(n)$ and there is nothing to be discussed). First, consider type $[\mathrm{S}H^{+}]$. For the notation, see (15). Let H be a Howe subgroup of $\mathrm{O}(n)$ in standard form and let Q be a principal $\mathrm{S}H^{+}$-bundle over X. Since the identity connected component H_0 of H is also the identity connected component of $\mathrm{S}H^{+}$ and since X is simply connected, any connected component of Q is a principal H_0-bundle. One may use the classification of H_0-bundles to describe Q in terms of the characteristic classes of its connected components. In general, the latter will not be isomorphic, so that one would have to identify their characteristic classes appropriately to obtain a unique description of Q. Although this can be done easily we will pursue another approach here.
For a Lie subgroup $G \subseteq G'$ and a principal G-bundle P over X, let $P^{G'}$ denote the extension of P to the structure group G'. This is the unique principal G'-bundle which contains P as a reduction to the subgroup G.[5] Consider the principal H-bundle Q^H obtained from Q by extension of the structure group to H. Q is a reduction of that bundle to the subgroup $\mathrm{S}H^{+}$. Since $H/\mathrm{S}H^{+} = \mathbb{Z}_2$, there exists exactly one further such reduction Q', which might be isomorphic to Q but in general is not. Thus, we may classify $\mathrm{S}H^{+}$-bundles by the characteristic classes of H-bundles and a certain $\mathbb{Z}_2$-valued quantity. To construct it, choose unique representatives in $H^*(X, \mathbb{Z})$ of the classes in $\mathrm{P}H^*(X, \mathbb{Z})$ and define a map $\sigma : H^*(X, \mathbb{Z}) \to \mathbb{Z}_2$ by

$$\sigma(\alpha) = \begin{cases} 1 & \alpha \text{ is a representative,} \\ -1 & \alpha \text{ is not.} \end{cases}$$

For a principal H_0-bundle Q_0 define

$$\sigma(Q_0) := \prod_{\mathbb{K}_i = \mathbb{R}} \sigma\big(e(Q_{0i})\big)^{k_i} .$$

Lemma 4.20. *Let Q_0 and Q_0' be principal H_0-bundles over X which have isomorphic extensions to H. Then Q_0 and Q_0' have isomorphic extensions to $\mathrm{S}H^{+}$ if and only if $\sigma(Q_0) = \sigma(Q_0')$.*

Proof. We may identify Q_0 and Q_0' with subsets of the extension $Q_0^H \equiv Q_0'^H = \tilde{Q}$. We claim that there exists $h \in H$ transforming Q_0 to Q_0' inside $\tilde{Q}$. To see this, consider the quotient bundle $\tilde{Q}/H_0$. This is a principal bundle over X with structure group H/H_0. The natural projection $p : \tilde{Q} \to \tilde{Q}/H_0$ together with the homomorphism $H \to H/H_0$ yields a principal bundle morphism covering the identity on X. There exist sections s, s' in $\tilde{Q}/H_0$ such that $Q_0 = p^{-1}(s(X))$ and $Q_0' = p^{-1}(s'(X))$. Since $\tilde{Q}/H_0$ has finite fibers, s and s' are determined by their values at a single point. Hence, there exists an element of H/H_0 transforming s to s'. Then any representative of this element transforms Q_0 to Q_0'.

---

[5] $P^{G'}$ can be constructed as the fiber bundle associated with P by virtue of the action of G on G' by left multiplication. The action of G' on $P^{G'}$ is induced from the action of G' on itself by right multiplication.

Now let h_i denote the projections of h to the factors of H. Then

$$\det(h) = \textstyle\prod_{\mathbb{K}_i=\mathbb{R}} \det(h_i)^{k_i} \,. \tag{35}$$

The group element h_i transforms the factor Q_{0i} of Q_0 to the factor Q'_{0i} of Q'_0, where both factors are viewed as subbundles of the corresponding factor of $\tilde{Q}$. Hence, for the factors with $\mathbb{K}_i = \mathbb{R}$ there holds

$$e(Q'_{0i}) = \det(h_i) e(Q_{0i}) \,. \tag{36}$$

In case $e(Q_{0i}) = -e(Q_{0i})$, the factors Q_{0i} and Q'_{0i} are isomorphic, no matter what determinant h_i has. By passing from Q'_0 to an isomorphic subbundle, denoted by the same symbol, we may assume that $h_i = \mathbb{1}$ in that case. Under this assumption on h, (35) and (36) yield

$$\sigma(Q'_0) = \det(h)\, \sigma(Q_0) \,. \tag{37}$$

If $\sigma(Q'_0) = \sigma(Q_0)$, (37) implies that $h \in \mathrm{S}H^+$ and hence $Q_0^{\mathrm{S}H^+} = Q_0'^{SH^+}$. Conversely, if $Q_0^{SH^+} = Q_0'^{SH^+}$ then Q_0 and Q'_0 project to the same section in the quotient bundle $\tilde{Q}/\mathrm{S}H^+$. Then h projects to the class of the identity in $H/\mathrm{S}H^+$, i.e., $h \in \mathrm{S}H^+$. Hence, (37) implies $\sigma(Q'_0) = \sigma(Q_0)$. □

Lemma 4.20 yields in particular that for an $\mathrm{S}H^+$-bundle Q we may define $\sigma(Q)$ to be $\sigma(Q_0)$ for some connected component Q_0. By construction, this definition applies when H possesses an O-factor. If it does not, we define $\sigma(Q) = 1$. Then Theorem 4.19 and Lemma 4.20 imply

Theorem 4.21. *Let X be a simply connected CW-complex of dimension 4 without 2-torsion in $H^4(X, \mathbb{Z})$ and let $H = \mathrm{I}_{\mathbb{K}_1}{}^{k_1}_{m_1} \times \cdots \times \mathrm{I}_{\mathbb{K}_r}{}^{k_r}_{m_r}$ be a Howe subgroup of $\mathrm{O}(n)$ of standard form. Principal $\mathrm{S}H^+$-bundles are classified by the characteristic classes of the factors of their extension to the structure group H and the invariant σ.* □

Finally, consider type $[\mathrm{S}H^-]$. For a principal bundle Q with structure group $\mathrm{S}H^-$ we form the extension $Q^{\mathrm{O}(n)}$ and transform Q inside $Q^{\mathrm{O}(n)}$ by the element $a_{(n)}$ of $\mathrm{O}(n)$ we had chosen for the definition of $\mathrm{S}H^-$. (We could take any $a \in \mathrm{O}(n)$ such that $a\mathrm{S}H^- a^{-1} = \mathrm{S}H^+$.) Since the transformed bundle has structure group $\mathrm{S}H^+$, we can define $\sigma(Q)$ and the characteristic classes of Q to be given by the corresponding quantities of the transformed bundle.

Remark 4.22. Since the Euler class is a 2-torsion element in odd dimension, only SO(2)- and SO(4)-factors contribute to σ. If moreover one is interested in holonomy-induced Howe subbundles only, SO(2)-factors do not appear, hence here it is just the Euler classes of the SO(4)-factors which determine σ.

We now pass to manifolds as base spaces. It was shown by Thom [42] and Milnor [32] that every closed n-dimensional manifold M is homeomorphic to a finite n-dimensional CW-complex. Thus, all the above classification results carry over to bundles over closed manifolds. Since in our situation M is in addition simply connected and hence orientable, Poincaré duality holds. This implies:

- $H^n(M, \mathbb{Z})$ is torsion-free. Hence, Theorems 4.16, 4.19 and 4.21 as well as Corollary 4.18 apply to closed and simply connected manifolds of dimension 4.
- $H^3(M, \mathbb{Z}) = 0$. Hence the Euler class of any SO(3)-bundle vanishes.
- $H^3(M, \mathbb{Z}_2) = 0$. Hence, the induced homomorphism θ_* is injective in dimension 4. Then $w_4(P)$ is uniquely determined by the fundamental relation (34) in terms of $w_2(P)$ and $p_1(P)$:

$$\theta_* w_4(P) = \mathfrak{P} w_2(P) - \rho_4 p_1(P) \,. \tag{38}$$

This means that we can ignore w_4 in the classification of $\mathrm{SO}(n)$- and $\mathrm{O}(n)$-bundles over M. Replacing w_4 by (38) in the relations listed in Theorem 4.16, we obtain new relations involving only w_2, p_1 and e or $[e]$, respectively, and the fundamental relation is automatically satisfied. In particular, for $\mathrm{O}(n)$ and $\mathrm{SO}(n)$ with $n \geq 5$ there are no relations any more.

Remark 4.23. If we drop the restriction on X to be simply connected, the bundles need not be orientable so that the classification of $[X, \mathrm{B}\,\mathrm{O}(4)]$ may not be reduced to the classification of $[X, \mathrm{B}\,\mathrm{SO}(4)]$ anymore. Presumabely the classification can be done for orientable four dimensional base complexes using similar methods as in the proof of Theorem 4.9, but replacing the ordinary Euler class by the Euler class with local coefficients.

5 Howe subbundles

In this section we determine the isomorphism classes of Howe subbundles of principal bundles P with structure group $G = \mathrm{O}(n)$, $\mathrm{SO}(n)$ or $\mathrm{Sp}(n)$ over closed simply connected manifolds of dimension 4. Since

- subsequently we have to further factorize the isomorphism classes by the action of the structure group G of P,
- this action transforms the structure group of the reduction by conjugation,
- any Howe subgroup is conjugate to one of standard form, see Section 3,

it suffices to discuss reductions of P to Howe subgroups of standard form. Such reductions will be referred to as Howe subbundles of standard form.

First, we explain the idea. If a principal H-bundle Q and a principal G-bundle P are given then Q is a reduction of P if and only if the extension Q^G of Q to the structure group G is isomorphic to P. According to the results of Section 4, in the case where $G = \mathrm{O}(n)$, $\mathrm{SO}(n)$ or $\mathrm{Sp}(n)$ and M is closed simply connected and of dimension 4 it suffices to compare the characteristic classes. Thus, all we have to do is to compute the characteristic classes of the G-bundle Q^G in terms of the characteristic classes of the H-bundle Q. The classifying map of Q^G is $\mathrm{B}\,j \circ f$, where $f : M \to \mathrm{B}\,H$ is the classifying map of Q and $\mathrm{B}\,j : \mathrm{B}\,H \to \mathrm{B}\,G$ is the classifying map of the extension of the universal H-bundle to the structure group G. Consequently, $\alpha(Q^G) = f^* \circ (Bj)^* \alpha$. It follows that Q is a reduction of P iff there holds $\alpha(P) = f^* \circ (Bj)^* \alpha$ for any characteristic class α of G-bundles. This provides a set of equations in the cohomology of M whose solutions classify the reductions of P to the structure group H.

5.1 Howe subbundles of principal $\mathrm{O}(n)$- and $\mathrm{SO}(n)$-bundles

We treat reductions of $\mathrm{O}(n)$ to Howe subgroups and reductions of $\mathrm{SO}(n)$ to subgroups of type $[\mathrm{S}H^{\pm}]$ simultaneously. Let $H = \mathrm{I}_{\mathbb{K}_1}{}^{k_1}_{m_1} \times \cdots \times \mathrm{I}_{\mathbb{K}_r}{}^{k_r}_{m_r}$ be a Howe subgroup of $\mathrm{O}(n)$. We observe: since M is simply connected, a principal H-bundle Q is a reduction of a principal $\mathrm{O}(n)$-bundle P iff there exists a common reduction of Q and P to H_0. A similar statement holds for H and $\mathrm{O}(n)$ replaced by $\mathrm{S}H^+$ and $\mathrm{SO}(n)$. Hence, it suffices to study the case of an H_0-bundle Q and to compute the characteristic classes of its extension $Q^{\mathrm{SO}(n)}$ to $\mathrm{SO}(n)$, i.e., $\alpha\big(Q^{\mathrm{SO}(n)}\big)$, where $\alpha = w,\ p,\ e$. Let $f_i : M \to \mathrm{B}\,\mathrm{I}_{\mathbb{K}_i}(m_i)_0$ denote the classifying map of the factor Q_i of Q. For any set X let $\Delta_r : X \to X \times \overset{r}{\cdots} \times X$ denote the diagonal map. For clarity, in the calculation to follow we label characteristic classes by the group they belong to.

The classifying map of Q is given by

$$f = \Big(\prod\nolimits_i f_i\Big) \circ \Delta_r\,. \tag{39}$$

The classifying map of $Q^{\mathrm{SO}(n)}$ is $\mathrm{B}\, j \circ f$, where $\mathrm{B}\, j : \mathrm{B}\, H_0 \to \mathrm{B}\, \mathrm{SO}(n)$ is the map of classifying spaces induced by j. Hence,

$$\alpha(Q^{\mathrm{SO}(n)}) = f^* \, (\mathrm{B}\, j)^* \alpha(\mathrm{SO}(n)) \,, \qquad \alpha = w, p, e \,. \tag{40}$$

We decompose $j : H_0 \to \mathrm{SO}(n)$ as

$$H_0 = \prod\nolimits_i \mathrm{I}_{\mathbb{K}_i}(m_i)_0 \xrightarrow{\prod_i \varphi_{\mathbb{K}_i,\mathbb{R}}} \prod\nolimits_i \mathrm{SO}(\delta_i m_i) \xrightarrow{\prod_i \Delta_{k_i}} \prod\nolimits_i \mathrm{SO}(\delta_i m_i) \times \overset{k_i}{\cdots} \times \mathrm{SO}(\delta_i m_i) \xrightarrow{i} \mathrm{SO}(n) \,,$$

where $\delta_i = \dim_{\mathbb{R}}(\mathbb{K}_i)$ and i denotes the standard blockwise embedding. Hence,

$$(\mathrm{B}\, j)^* \; \alpha(\mathrm{SO}(n)) = \Big(\prod\nolimits_i \big(\mathrm{B}\, \varphi_{\mathbb{K}_i,\mathbb{R}} \big)^* \Big) \circ \Big(\prod\nolimits_i \Delta_{k_i}^* \Big) \circ (\mathrm{B}\, i)^* \; \alpha\big(\mathrm{SO}(n) \big) \,. \tag{41}$$

Since for the standard embedding $\varphi : \mathrm{SO}(l) \to \mathrm{SO}(l+1)$ there holds $(\mathrm{B}\, \varphi)^* \alpha(\mathrm{SO}(l+1)) = \alpha(\mathrm{SO}(l))$,

$$(\mathrm{B}\, i)^* \alpha(\mathrm{SO}(n)) = \prod\nolimits_i \alpha(\mathrm{SO}(\delta_i m_i)) \times \overset{k_i}{\cdots} \times \alpha(\mathrm{SO}(\delta_i m_i)) \,, \tag{42}$$

where $\times$ and $\prod_i$ refer to the cohomology cross product. Eqs. (39), (40), (41) and (42) imply

$$\alpha\left(Q^{\mathrm{SO}(n)} \right) = \Big(f_1^* \, \big(\mathrm{B}\, \varphi_{\mathbb{K}_1,\mathbb{R}} \big)^* \alpha(\mathrm{SO}(\delta_1 m_1)) \Big)^{k_1} \cdots \Big(f_r^* \, \big(\mathrm{B}\, \varphi_{\mathbb{K}_r,\mathbb{R}} \big)^* \alpha(\mathrm{SO}(\delta_r m_r)) \Big)^{k_r} \,, \tag{43}$$

where powers and products refer to the cup product on M. Since $\mathrm{B}\, \varphi_{\mathbb{K}_i,\mathbb{R}} \circ f_i$ is the classifying map of the extension $\hat{Q}_i := Q_i^{\mathrm{SO}(\delta_i m_i)}$ of the factor Q_i ('realification'), we can simplify (43) to

$$\alpha\left(Q^{\mathrm{SO}(n)} \right) = \alpha\big(\hat{Q}_1\big)^{k_1} \cdots \alpha\big(\hat{Q}_r\big)^{k_r} \,. \tag{44}$$

For $\mathbb{K}_i = \mathbb{R}$, we have $\delta_i = 1$ and $\alpha\big(\hat{Q}_i\big) = \alpha\big(Q_i\big)$. For $\mathbb{K}_i = \mathbb{C}, \mathbb{H}$, denote $\tilde{p} = \sum_k (-1)^k p_k$ and $\tilde{c} = \sum_k (-1)^k c_k$. The standard relations between Chern classes and real characteristic classes listed in Proposition D.5 imply

$$w\big(\hat{Q}_i\big) = \rho_2 \, c\big(Q_i\big) \,, \qquad \tilde{p}\big(\hat{Q}_i\big) = c\big(Q_i\big) \tilde{c}\big(Q_i\big) \,, \qquad e\big(\hat{Q}_i\big) = c_{\frac{1}{2}\delta_i m_i}\big(Q_i\big) \,. \tag{45}$$

Theorem 5.1. *Let* $H = \mathrm{I}_{\mathbb{K}_1}{}^{k_1}_{m_1} \times \cdots \times \mathrm{I}_{\mathbb{K}_r}{}^{k_r}_{m_r}$ *be a Howe subgroup of* $\mathrm{O}(n)$ *and let* M *be a closed simply connected* 4*-manifold. Let* P *be a principal* $\mathrm{SO}(n)$*-bundle over* M *and let* Q *be a principal* H_0*-bundle over* M*. Then* Q *is a reduction of* P *if and only if*

$$\begin{aligned}
w(P) &= \prod\nolimits_{\mathbb{K}_i = \mathbb{R}} \; w(Q_i)^{k_i} \cdot \prod\nolimits_{\mathbb{K}_i = \mathbb{C},\mathbb{H}} \; \rho_2 \left(c(Q_i) \right)^{k_i} \,, && (46)\\
\tilde{p}(P) &= \prod\nolimits_{\mathbb{K}_i = \mathbb{R}} \; \tilde{p}(Q_i)^{k_i} \cdot \prod\nolimits_{\mathbb{K}_i = \mathbb{C},\mathbb{H}} \; c(Q_i)^{k_i} \, \tilde{c}(Q_i)^{k_i} \,, && (47)\\
e(P) &= \prod\nolimits_{\mathbb{K}_i = \mathbb{R}} \; e(Q_i)^{k_i} \cdot \prod\nolimits_{\mathbb{K}_i = \mathbb{C}} \; c_{m_i}(Q_i)^{k_i} \cdot \prod\nolimits_{\mathbb{K}_i = \mathbb{H}} \; c_{2 m_i}(Q_i)^{k_i} \,, && (48)
\end{aligned}$$

Proof. As noted above, Q is a reduction of P iff $Q^{\mathrm{SO}(n)}$ is isomorphic to P, hence iff $\alpha(Q^{\mathrm{SO}(n)}) = \alpha(P)$ for $\alpha = w$, p and e. Equations (46)–(48) then follow from Equations (44) and (45), where for (47) one has to use that $p \mapsto \tilde{p}$ commutes with the cup product. ☐

By the arguments given above, Theorem 5.1 solves the problem of Howe subbundles of principal bundles with structure groups $\mathrm{O}(n)$ and $\mathrm{SO}(n)$. It remains to rewrite the reduction equations (46)–(48) in terms of the quantities which classify H-bundles and $\mathrm{S}H^{\pm}$-bundles, respectively.

Corollary 5.2. *Under the assumptions of Theorem* 5.1*, let* P *be a principal* $\mathrm{O}(n)$*-bundle over* M *and let* Q *be a principal* H*-bundle over* M*. Then* Q *is a reduction of* P *if and only if Equations* (46) *and* (47) *hold, as well as*

$$[e](P) \quad = \quad \prod\nolimits_{\mathbb{K}_i=\mathbb{R}} [e](Q_i)^{k_i} \cdot \prod\nolimits_{\mathbb{K}_i=\mathbb{C}} [c_{m_i}(Q_i)]^{k_i} \cdot \prod\nolimits_{\mathbb{K}_i=\mathbb{H}} [c_{2m_i}(Q_i)]^{k_i} \,, \tag{49}$$

where taking the class $[c_{m_i}(Q_i)]$ *means that* $c_{m_i}(Q_i)$ *is identified with* $-c_{m_i}(Q_i)$*.*

Proof. As noted above, Q is a reduction of P iff there exists a common reduction Q_0 to H_0. This is equivalent to the condition that there exist reductions Q_0 of Q to H_0 and P_0 of P to $\mathrm{SO}(n)$ such that Q_0 is a reduction of P_0, i.e., such that Equations (46)–(48) hold with P replaced by P_0 and Q_i replaced by Q_{0i}. By using that for $\mathbb{K}_i = \mathbb{C}$ and $\mathbb{H}$ one has $Q_{0i} = Q_i$, in (46) and (47) we can then replace P_0 by P and Q_{0i} by Q_i again, with Q and P now meaning the bundles given here. Equation (49) follows from (48) using that the cup product commutes with passing to equivalence classes w.r.t. the equivalence relation $\alpha \sim \beta \Leftrightarrow \alpha + \beta = 0$. ☐

Corollary 5.3. *Under the assumptions of Theorem* 5.1*, let* P *be a principal* $\mathrm{SO}(n)$*-bundle over* M *and let* Q *be a principal* $\mathrm{S}H^{\pm}$*-bundle over* M*. Then* Q *is a reduction of* P *if and only if for the factors* Q_i *of the extension of* Q *to the structure group* H *there hold Eqs.* (46) *and* (47)*, as well as*

$$e(P) \quad = \quad \pm\sigma(Q) \cdot \prod\nolimits_{\mathbb{K}_i=\mathbb{R}} e^{\circ}(Q_i)^{k_i} \cdot \prod\nolimits_{\mathbb{K}_i=\mathbb{C}} c_{m_i}(Q_i)^{k_i} \cdot \prod\nolimits_{\mathbb{K}_i=\mathbb{H}} c_{2m_i}(Q_i)^{k_i} \,, \tag{50}$$

where the postitive sign applies to $\mathrm{S}H^{+}$ *and the negative sign to* $\mathrm{S}H^{-}$ *and where* $e^{\circ}(Q_i)$ *denotes the representative for the equivalence class* $[e](Q_i)$ *chosen in the definition of* σ*.*

Proof. If Q is an $\mathrm{S}H^{+}$-bundle, it is a reduction of P iff there exists a common reduction Q_0 to H_0, i.e., iff there exists an H_0-bundle such that (46)–(48) hold with Q_i replaced by Q_{0i}. By the same argument as in the proof of Corollary 5.2, in (46) and (47) we can replace Q_{0i} by Q_i again, with Q_i now meaning the factors of the extension Q^{H}. Equation (50) follows from (48) by replacing $e(Q_{0i}) = \sigma\big(e(Q_{0i})\big)\, e^{\circ}(Q_i)$. If Q is an $\mathrm{S}H^{-}$-bundle, we pass to $Q^{\mathrm{O}(n)}$ and $P^{\mathrm{O}(n)}$ and transform both Q inside $Q^{\mathrm{O}(n)}$ and P inside $P^{\mathrm{O}(n)}$ by the element $a_{(n)}$ of $\mathrm{O}(n)$ chosen to define $\mathrm{S}H^{-}$. Let the transforms be denoted by $\tilde{Q}$ and $\tilde{P}$, respectively. We have $Q \subseteq P$ iff $\tilde{Q} \subseteq \tilde{P}$, i.e., iff (46), (47) and (50) hold with Q and P replaced by $\tilde{Q}$ and $\tilde{P}$, respectively. By definition of the classifying data of Q, in these equations we can replace $\tilde{Q}$ by Q again. Since $w(\tilde{P}) = w(P)$, $p(\tilde{P}) = p(P)$ and $e(\tilde{P}) = -e(P)$, the assertion follows. ☐

Remark 5.4. The powers of characteristic classes appearing in Corollaries 5.2 and 5.3 amount to

$$\begin{aligned}
w(Q_i)^{k_i} &= 1 + k_i w_2(Q_i) + k_i w_3(Q_i) + k_i w_4(Q_i) + \tbinom{k_i}{2} w_2(Q_i)^2 \,, \\
c(Q_i)^{k_i} &= 1 + k_i c_1(Q_i) + k_i c_2(Q_i) + \tbinom{k_i}{2} c_1(Q_i)^2 \,, \\
\tilde{p}(Q_i)^{k_i} &= 1 - k_i p_1(Q_i) \,, \\
c(Q_i)^{k_i} \tilde{c}(Q_i)^{k_i} &= 1 + 2k_i c_2(Q_i) - k_i c_1(Q_i)^2 \,.
\end{aligned}$$

We observe:

– Equation (46) has to be analyzed in dimension 2 only;

– Equations (48), (49) and (50) have to be analyzed for $n \neq 2, 4$ only.

Indeed, due to (31), if (46) holds in dimension 2 then it holds in dimension 3. Similarly, if (47) is satisfied and if (46) holds in dimension 2 then it is automatically satisfied in dimension 4, because $w_4(P)$ is determined by $w_2(P)$ and $p_1(P)$, see 38. Finally, for $n \neq 2, 4$, Equations (48), (49) and (50) are automatically satisfied, because for $n = 3$, H has an O-factor of odd rank and for $n \geq 5$ the product of Euler classes vanishes identically because of the dimension.

Remark 5.5. The reduction equations (46)–(48), (49) or (50) are supplemented by the relations between the characteristic classes of the SO- or O-factors, respectively, listed in Theorem 4.16 or Corollary 4.18, respectively. As observed above, in these relations, $w_4(Q_i)$ can be replaced by $w_2(Q_i)$ and $p_1(Q_i)$ using (38).

5.2 Howe subbundles of principal $\mathrm{Sp}(n)$-bundles

Let a Howe subgroup $H = \mathrm{I}_{\mathbb{K}_1}{}^{k_1}_{m_1} \times \cdots \times \mathrm{I}_{\mathbb{K}_r}{}^{k_r}_{m_r}$ of $\mathrm{Sp}(n)$ and a principal H-bundle Q be given. We have to compute $c\big(Q^{\mathrm{Sp}(n)}\big)$. The classifying map of Q is given by $f = (\prod_i f_i) \circ \Delta_r$, where $f_i : M \to \mathrm{B}\,\mathrm{I}_{\mathbb{K}_i}(m_i)$ are the classifying maps of the factors Q_i of Q. The classifying map of $Q^{\mathrm{Sp}(n)}$ is $\mathrm{B}\,j \circ f$, where $j : H \to \mathrm{Sp}(n)$ denotes the natural embedding. Hence

$$c\big(Q^{\mathrm{Sp}(n)}\big) = f^*(\mathrm{B}\,j)^* c\big(\mathrm{Sp}(n)\big)\,.$$

We decompose j as

$$H = \prod\nolimits_i \mathrm{I}_{\mathbb{K}_i}(m_i) \xrightarrow{\prod_i j_{\mathbb{K}_i,\mathbb{H}}} \prod\nolimits_i \mathrm{Sp}(m_i) \xrightarrow{\prod_i \Delta_{k_i}} \prod\nolimits_i \mathrm{Sp}(m_i) \times \overset{k_i}{\cdots} \times \mathrm{Sp}(m_i) \xrightarrow{i} \mathrm{Sp}(n)\,,$$

where $j_{\mathbb{K}_i,\mathbb{H}} : \mathrm{I}_{\mathbb{K}_i}(m_i) \to \mathrm{Sp}(m_i)$ denotes the natural embedding as a subset. Argueing as in Subsection 5.1 we arrive at

$$c\left(Q^{\mathrm{Sp}(n)}\right) = \left(f_1^*\,\left(\mathrm{B}\,j_{\mathbb{K}_1,\mathbb{H}}\right)^* c(\mathrm{Sp}(m_1))\right)^{k_1} \cdots \left(f_r^*\,\left(\mathrm{B}\,j_{\mathbb{K}_r,\mathbb{H}}\right)^* c(\mathrm{Sp}(m_r))\right)^{k_r}. \tag{51}$$

Since $\mathrm{B}\,j_{\mathbb{K}_i,\mathbb{H}} \circ f_i$ is the classifying map of the extension $Q_i^{\mathrm{Sp}(m_i)}$, (51) can be written as

$$c\left(Q^{\mathrm{Sp}(n)}\right) = c\big(Q_1^{\mathrm{Sp}(m_1)}\big)^{k_1} \cdots c\big(Q_r^{\mathrm{Sp}(m_r)}\big)^{k_r}. \tag{52}$$

For $\mathbb{K}_i = \mathbb{H}$ one has $Q_i^{\mathrm{Sp}(m_i)} = Q_i$ and hence $c\big(Q_i^{\mathrm{Sp}(m_i)}\big) = c(Q_i)$. The standard relations between characteristic classes listed in Proposition D.5 yield

$$c\big(Q_i^{\mathrm{Sp}(m_i)}\big) = \begin{cases} \tilde{p}(Q_i)^2 & \mathbb{K}_i = \mathbb{R}\,, \\ c(Q_i)\,\tilde{c}(Q_i) & \mathbb{K}_i = \mathbb{C}\,. \end{cases}$$

Combining this with (52) we arrive at

Theorem 5.6. *Let $H = \mathrm{I}_{\mathbb{K}_1}{}^{k_1}_{m_1} \times \cdots \mathrm{I}_{\mathbb{K}_r}{}^{k_r}_{m_r}$ be a Howe subgroup of $\mathrm{Sp}(n)$ and let M be a closed simply connected 4-manifold. Let P be a principal $\mathrm{Sp}(n)$-bundle over M and let Q be a principal H-bundle over M. Then Q is a reduction of P if and only if*

$$c(P) = \prod\nolimits_{\mathbb{K}_i=\mathbb{R}} \tilde{p}(Q_i)^{2k_i} \cdot \prod\nolimits_{\mathbb{K}_i=\mathbb{C}} c(Q_i)^{k_i}\,\tilde{c}(Q_i)^{k_i} \cdot \prod\nolimits_{\mathbb{K}_i=\mathbb{H}} c(Q_i)^{k_i}\,. \tag{53}$$

□

For explicit formulae for the powers appearing in (53) see Remark 5.4. Like in the case of $\mathrm{O}(n)$-bundles and $\mathrm{SO}(n)$-bundles, the reduction equation is supplemented by the relations between the characteristic classes of O-factors, see Remark 5.5 for details.

6 Holonomy-induced Howe subbundles

According to our programme, next we have to specify the Howe subbundles which are holonomy-induced, i.e., generated by a connected subbundle. Over a simply connected manifold, this turns out to be a purely algebraic condition.

Lemma 6.1. *Let P be a principal G-bundle over M. If M is simply connected then a reduction of P to a Howe subgroup H of G is holonomy-induced if and only if H is generated by its identity connected component H_0. That is to say, $H = \mathrm{C}^2_G(H_0)$ or, alternatively, $\mathrm{C}_G(H) = \mathrm{C}_G(H_0)$.*

Proof. Let H be given and let Q be a reduction of P to H. First, assume that $H = \mathrm{C}^2_G(H_0)$. Since M is simply connected, Q can be reduced to H_0. By assumption, Q is the Howe subbundle generated by the reduction. Since H_0 is connected, the reduction is connected[6]. Conversely, assume that Q is holonomy-induced. Let Q_1 be a connected subbundle generating Q as a Howe subbundle and let H_1 be the structure group of Q_1. Then $H = \mathrm{C}^2_G(H_1)$. We show that H_1 is connected, hence $H_1 = H_0$. For that purpose, it suffices to show that the fibers of Q_1 are connected. Thus, let $q_a, q_b \in Q_1$ with common base point $m \in M$. Since Q_1 is connected, there exists a path γ from q_a to q_b. The path γ projects to a closed path in M based at m. Since M is simply connected, the projected path is homotopic to the constant path through m. Consequently, by the homotopy lifting property, there exists a map $H : [0,1] \times [0,1] \to Q_1$ such that $H(0,\cdot) = \gamma$ and $H(1,\cdot)$ is a path in the fiber of Q_1 over m. Then the three pieces $H(\cdot,0)$, $H(1,\cdot)$ and $H(1-\cdot,1)$ establish a path from q_a to q_b inside this fiber. □

Thus, we have to determine the Howe subgroups satisfying

$$H = \mathrm{C}^2_G(H_0) \quad \Leftrightarrow \quad \mathrm{C}_G(H) = \mathrm{C}_G(H_0)\,. \tag{54}$$

First, we will treat the cases $G = \mathrm{O}(n)$ and $G = \mathrm{Sp}(n)$.

Lemma 6.2. *A Howe subgroup of a compact Lie group satisfies* (54) *if and only if it does not contain another Howe subgroup of the same dimension.*

Proof. Let G be compact and let H be a Howe subgroup. The subgroup $\mathrm{C}^2_G(H_0)$ is a Howe subgroup. Since $H_0 \subseteq \mathrm{C}^2_G(H_0) \subseteq H$, it has the same dimension as H. Hence, if H does not contain another Howe subgroup of the same dimension, (54) holds. Conversely, let $K \subset H$ be another Howe subgroup of the same dimension. Since K_0 and H_0 are closed (i.e. compact without boundary) connected manifolds of the same dimension, invariance of domain (see [29, p. 217, Ex. 6.5]) implies that the embedding $K_0 \subseteq H_0$ is an open map. Thus K_0 is an open and closed subset of H_0 and hence there holds $K_0 = H_0$. It follows $\mathrm{C}^2_G(H_0) = \mathrm{C}^2_G(K_0) \subseteq K \neq H$. □

Lemma 6.3. *A Howe subgroup of* $\mathrm{O}(n)$ *or* $\mathrm{Sp}(n)$ *contains another Howe subgroup of the same dimension if and only if it contains a factor* $\mathrm{O}(2)$ *or a double factor* $\mathrm{O}(1) \times \mathrm{O}(1)$.

Proof. We give the argument for $G = \mathrm{O}(n)$; the case of $\mathrm{Sp}(n)$ is completely analogous. Consider the operations producing direct predecessors in the set of Howe subgroups of $\mathrm{O}(n)$, listed in Section 3.1. A given Howe subgroup H contains another Howe subgroup of the same dimension iff an operation can be applied to H which does not change the dimension. As noted in the proof of Lemma 3.4, the operations which do not change the dimension are inverse field restriction of a factor $\mathrm{O}(2)$ and merging of a double factor $\mathrm{O}(1) \times \mathrm{O}(1)$. ☐

---

[6] The implication under consideration holds without the assumption that M be simply connected; using this assumption simplifies the proof though.

Lemmas 6.1–6.3 imply

Theorem 6.4. *A Howe subbundle of a principal* $\mathrm{O}(n)$*- or* $\mathrm{Sp}(n)$*-bundle is holonomy-induced if and only if its structure group does not contain a factor* O_2^k *nor a double factor* $\mathrm{O}_1^k \times \mathrm{O}_1^l$. ☐

Next, we will treat the case of $G = \mathrm{SO}(n)$. Since in the case of odd n the argument is not simpler than in the case of even n and since the description of Howe subgroups of $\mathrm{SO}(n)$, n even, in terms of S-admissible Howe subgroups of $\mathrm{O}(n)$ trivially applies to the case of odd n, too, we can treat both cases at once, always working with S-admissible Howe subgroups of $\mathrm{O}(n)$.

Lemma 6.5. *Let H be an* S*-admissible Howe subgroup of* $\mathrm{O}(n)$*. Then* $\mathrm{S}H$ *contains another Howe subgroup of* $\mathrm{SO}(n)$ *of the same dimension if and only if H contains another Howe subgroup of* $\mathrm{O}(n)$ *of the same dimension.*

Proof. If H does not contain another Howe subgroup of $\mathrm{O}(n)$ of the same dimension, then H satisfies $\mathrm{C}_{\mathrm{O}(n)}(H) = \mathrm{C}_{\mathrm{O}(n)}(H_0)$ by Lemma 6.2. Since $H_0 \subseteq \mathrm{S}H \subseteq H$, there holds $\mathrm{C}_{\mathrm{O}(n)}(H_0) \supseteq \mathrm{C}_{\mathrm{O}(n)}(\mathrm{S}H) \supseteq \mathrm{C}_{\mathrm{O}(n)}(H)$, hence $C_{\mathrm{O}(n)}(H_0) = \mathrm{C}_{\mathrm{O}(n)}(\mathrm{S}H)$. Intersecting with $\mathrm{SO}(n)$ and using $(\mathrm{S}H)_0 = H_0$ we obtain $\mathrm{C}_{\mathrm{SO}(n)}(\mathrm{S}H) = \mathrm{C}_{\mathrm{SO}(n)}\big((\mathrm{S}H)_0\big)$.
Conversely, assume that H contains another Howe subgroup K of $\mathrm{O}(n)$ of the same dimension. Then $\mathrm{S}K \subseteq \mathrm{S}H$. Since intersection with $\mathrm{SO}(n)$ does not change the dimension, $\mathrm{S}K$ has the same dimension as $\mathrm{S}H$. It remains to show that $\mathrm{S}K$ is a Howe subgroup of $\mathrm{SO}(n)$ and that it is distinct from $\mathrm{S}H$. For that purpose, it suffices to show that K is S-admissible. We may assume that K is a direct predecessor of H and thus is obtained from H by inverse field restriction of a factor O_2^k or by merging a double factor $\mathrm{O}_1^k \times \mathrm{O}_1^l$. We check that neither of these two operations produces a factor that causes K or $\mathrm{C}_{\mathrm{O}(n)}(K)$ to belong to cases (A) or (B) of Lemma 3.4. For inverse field restriction of a factor O_2^k this is obvious, as it produces a factor U_1^k. Merging a double factor $\mathrm{O}_1^k \times \mathrm{O}_1^l$ produces the factor O_1^{k+l}. This cannot make K belong to case (A) nor $\mathrm{C}_{\mathrm{O}(n)}(K)$ to case (B) but it might make K belong to case (B) or $\mathrm{C}_{\mathrm{O}(n)}(K)$ to case (A). If K belongs to case (B), $k + l$ is odd, hence exactly one of the two factors that are merged has odd multiplicity. Moreover, among the remaining factors of K there must be a single $\mathrm{O}(1)$-factor of odd multiplicity and no further O-factor of odd multiplicity. Since all the remaining factors are also present in H, H belongs to case (B) itself (contradiction). If $\mathrm{C}_{\mathrm{O}(n)}(K)$ belongs to case (A), there holds $k + l = 2$, hence $k = l = 1$, and K has no further O-factor of odd rank. Then $\mathrm{C}_{\mathrm{O}(n)}(H)$ belongs to case (B), which is a contradiction, too. ☐

Lemmas 6.1–6.3 and 6.5 imply

Theorem 6.6. *A Howe subbundle Q of a principal* $\mathrm{SO}(n)$*-bundle is holonomy-induced if and only if the Howe subgroup of* $\mathrm{O}(n)$ *generated by the structure group of Q does not contain a factor* O_2^k *nor a double factor* $\mathrm{O}_1^k \times \mathrm{O}_1^l$. ☐

When n is even, the conditions given in the theorem supersede conditions (A) and (B) of Lemma 3.4. Hence, in this case we arrive at the following characterization. The holonomy-induced Howe subbundles of a principal $\mathrm{SO}(n)$-bundle, n even, are given by the reductions to the subgroups $\mathrm{S}H$, where H is a Howe subgroup of $\mathrm{O}(n)$ which does not belong to one of the following cases:

- H contains a factor O_2^k,
- H contains a double factor $\mathrm{O}_1^k \times \mathrm{O}_1^l$,

- H contains a factor O^2_m with m odd and no further O-factor of odd rank,
- H contains a double factor $\mathrm{O}^1_k \times \mathrm{O}^1_l$ with k and l odd and no further O-factor of odd rank.

For illustration, Figures 4 and 5 show the Hasse diagrams of the sets of conjugacy classes of Howe subgroups satisfying (54) for some low-dimensional classical compact Lie groups. These are the structure groups of holonomy-induced Howe subbundles in the respective situations. The diagrams can be extracted from the Hasse diagrams in Figures 1, 2 and 3 by removing vertices which correspond to Howe subgroups containing a factor O^k_2 or a double factor $\mathrm{O}^k_1 \times \mathrm{O}^l_1$.

7 Factorization by the action of the structure group

Let P be a principal bundle with structure group $G = \mathrm{O}(n)$, $\mathrm{SO}(n)$ or $\mathrm{Sp}(n)$. In the last step of our program we have to factorize the set of isomorphism classes of holonomy-induced Howe subbundles of P by the action of the structure group G on subbundles. Since above we have already restricted attention to Howe subbundles of standard form, all we have to do now is to analyze which of these Howe subbundles of standard form are conjugate under the action of the structure group. Thus, below we will assume that we are given a reduction Q of P to some Howe subgroup H of G of standard form and determine which bundles arise from Q by transformation with $d \in G$ such that dHd^{-1} is of standard form again.

7.1 Structure groups $\mathrm{O}(n)$ and $\mathrm{Sp}(n)$

First, consider $G = \mathrm{O}(n)$ and $\mathrm{Sp}(n)$. Up to isomorphy, Howe subbundles of P of standard form are characterized by r-tuples

$$\big((\mathbb{K}_1, m_1, k_1, \alpha_1), \ldots, (\mathbb{K}_r, m_r, k_r, \alpha_r)\big) \tag{55}$$

where

- r is a positive integer,
- $\mathrm{I}_{\mathbb{K}_1}{}^{k_1}_{m_1} \times \cdots \times \mathrm{I}_{\mathbb{K}_r}{}^{k_r}_{m_r}$ is a Howe subgroup of G,
- α_i are characteristic classes of $\mathrm{I}_{\mathbb{K}_i}(m_i)$-bundles over M, satisfying the respective reduction equations derived in Section 5. That is, $\alpha_i = w, p, [e]$ for O-factors and $\alpha_i = c$ for U- and Sp-factors.

To derive how the structure group acts on these data we also have to consider the connected components of Howe subbundles of standard form. These are characterized analogously by the data (55), where now α_i are characteristic classes of $\mathrm{I}_{\mathbb{K}_i}(m_i)_0$-bundles over M. I.e., $\alpha_i = w, p, e$ for SO-factors and $\alpha_i = c$ for U- and Sp-factors.

We define the flip to be the following operation on factors. For an SO-factor, multiply the Euler class e by -1. For a U-factor, multiply the first Chern class c_1 by -1, i.e., replace c by $\tilde{c}$. For O-factors and Sp-factors, the flip is not defined.

Theorem 7.1. *Let P be a principal bundle with structure group $G = \mathrm{O}(n)$ or $\mathrm{Sp}(n)$ over a closed simply connected manifold of dimension 4. Two Howe subbundles of P of standard form are conjugate under the action of the structure group if and only if they can be transformed into one another by a permutation of factors and by flips.*

Proof. First, let $G = \mathrm{O}(n)$. Let there be given a Howe subbundle of P of standard form and let H denote the corresponding Howe subgroup. Let $d \in \mathrm{O}(n)$ such that dHd^{-1} is of standard form again. We have to find out how d acts on the data of a connected component Q_0 of Q and derive from that how it acts on the data of Q. Since dHd^{-1} is of standard form, we can



Figure 4: Hasse diagrams of the sets of conjugacy classes of Howe subgroups which are generated by their identity connected components for O(n), $n = 2, \ldots, 6$ and SO(n), $n = 3, \ldots, 6$. For the notation, see Figure 1.


O_1^1
U_1^1
Sp_1^1
$\mathrm{O}_1^1\,\mathrm{U}_1^1$
$\mathrm{O}_1^1\,\mathrm{Sp}_1^1$
$\mathrm{U}_1^1\,\mathrm{Sp}_1^1$
$\mathrm{Sp}_1^1\,\mathrm{Sp}_1^1$
O_1^2
U_1^2
$\mathrm{U}_1^1\,\mathrm{U}_1^1$
U_2^1
Sp_2^1
Sp_1^2

Figure 5: Hasse diagrams of the sets of conjugacy classes of Howe subgroups which are generated by their identity connected components for $G = \mathrm{Sp}(1)$ and $\mathrm{Sp}(2)$. For the notation, see Figure 1. For identifications due to $\mathrm{Sp}(1) \equiv \mathrm{SU}(2)$, see Figure 2.

decompose $d = d_1 d_2 d_3$, where d_1 is a pure permutation of factors, $d_2 \in \mathrm{C}_{\mathrm{O}(n)}(H)$ and d_3 leaves invariant each factor separately. The action of d_1 on Q is by the corresponding permutation of factors. The action of d_2 is trivial. The group element d_3 can be further decomposed as $d_3 = d_{31} \oplus \cdots \oplus d_{3r}$ with $d_{3i} \in \mathrm{N}_{\mathrm{O}(\delta_i m_i k_i)}\big(\mathrm{I}_{\mathbb{K}_i}{}^{k_i}_{m_i}\big)$. Each d_{3i} defines an automorphism φ_i of the identity connected component $\mathrm{I}_{\mathbb{K}_i}(m_i)_0$ of $\mathrm{I}_{\mathbb{K}_i}(m_i)$. According to Lemma 3.9, up to a redefinition of d_2 we may assume that $d_{3i} = a_i \oplus \cdots \oplus a_i$ (k_i times) for some $a_i \in \mathrm{N}_{\mathrm{O}(\delta_i m_i)}\big(\mathrm{I}_{\mathbb{K}_i}{}^{1}_{m_i}\big)$. Then φ_i is induced from the action of a_i on the subgroup $\big(\mathrm{I}_{\mathbb{K}_i}{}^{1}_{m_i}\big)_0$ of $\mathrm{O}(\delta_i m_i)$ by conjugation. We have to determine how $(\mathrm{B}\,\varphi_i)^*$ maps the characteristic classes of $\mathrm{I}_{\mathbb{K}_i}(m_i)_0$. If φ_i is inner, $(\mathrm{B}\,\varphi_i)^*\alpha_i = \alpha_i$, because $\mathrm{I}_{\mathbb{K}_i}(m_i)_0$ is arcwise connected. In the following situations, φ_i is outer, cf. Lemma 3.10:

- $\mathbb{K}_i = \mathbb{R}$, m_i even, $\det a_i = -1$. Here $(\mathrm{B}\,\varphi_i)^* w(\mathrm{SO}(m_i)) = w(\mathrm{SO}(m_i))$, $(\mathrm{B}\,\varphi_i)^* p(\mathrm{SO}(m_i)) = p(\mathrm{SO}(m_i))$ and $(\mathrm{B}\,\varphi_i)^* e(\mathrm{SO}(m_i)) = -e(\mathrm{SO}(m_i))$.
- $\mathbb{K}_i = \mathbb{C}$, $a_i = \mathbb{1}_{m_i,m_i}$. Here $(\mathrm{B}\,\varphi_i)^* c(\mathrm{U}(m_i)) = \tilde{c}(\mathrm{U}(m_i))$.

In either case, $(\mathrm{B}\,\varphi_i)^*$ induces a flip of the factor under consideration. Conversely, since a flip of an SO-factor of odd rank is trivial, any flip of Q_0 can be generated in this way. Finally, when passing to Q, flips of SO-factors cancel whereas flips of U-factors persist.

For $G = \mathrm{Sp}(n)$, the proof is analogous. Here, the automorphisms φ_i are induced by the action of $d_{3i} \in \mathrm{N}_{\mathrm{Sp}(m_i k_i)}\big(\mathrm{I}_{\mathbb{K}_i}{}^{k_i}_{m_i}\big)$ on the subgroup $\mathrm{I}_{\mathbb{K}_i}{}^{k_i}_{m_i}$ of $\mathrm{Sp}(m_i k_i)$. A case by case inspection of the outer automorphisms of $\mathrm{O}(m)$, $\mathrm{U}(m)$ and $\mathrm{Sp}(m)$ shows that here the φ_i are outer in the following situations:

- $\mathbb{K}_i = \mathbb{R}$, m_i even, $d_{3i} = a_i \oplus \cdots \oplus a_i$ (k_i times) with $a_i \in \mathrm{O}(m_i)$ such that $\det a_i = -1$;
- $\mathbb{K}_i = \mathbb{C}$, $d_{3i} = \mathrm{j}\mathbb{1}_{m_i k_i}$; here j denotes the second quaternionic unit.

The action of $(\mathrm{B}\varphi_i)^*$ on the characteristic classes is the same as for $G = \mathrm{O}(n)$. Hence, the assertion. □

7.2 Structure group SO(n)

Let P be a principal $\mathrm{SO}(n)$-bundle. As before, we will more generally consider reductions to subgroups of type $[\mathrm{S}H^{\pm}]$, see (15) and the comment after this formula. Such reductions will be referred to as reductions of type $[\mathrm{S}H^{\pm}]$. Isomorphy classes of such reductions are characterized

by the data (55) for Howe subbundles of $P^{\mathrm{O}(n)}$, together with a compatible value of the invariant σ. Like in the case of $G = \mathrm{O}(n)$, to prove the result for the reductions of type $[\mathrm{S}H^{\pm}]$ we have to consider their connected components, i.e., the reductions of P to the identity connected components of Howe subgroups of $\mathrm{O}(n)$. These are characterized by the same data as for $G = \mathrm{O}(n)$ where now the reduction equations are those of the $\mathrm{SO}(n)$-bundle P.
To formulate the result we need some more terminology. SO-factors of odd multiplicity, O-factors of odd multiplicity and U-factors of odd rank and odd multiplicity will be called signed factors. All other factors are called unsigned. A flip is called signed or unsigned according to the factor it is applied to. An SO-factor or a U-factor is called stable if it does not change under a flip, i.e., if for its characteristic class(es) there holds $-e = e$ or $\tilde{c} = c$, respectively. Otherwise, it is called unstable. An O-factors is called stable (unstable) if it arises from a stable (unstable) SO-factor by extension. This definition makes sense, because the SO-factor in question is determined by the given O-factor up to the sign of the Euler class. Note that to be signed is an algebraic property, depending on the structure group of the bundle reduction under consideration, whereas to be stable is a topological property, depending on the bundle reduction itself. Finally, if σ can take the value -1, there exists an unstable signed O-factor. In this case, we define the flip of σ as the passage to $-\sigma$. We assume this flip to be signed. Thus, unsigned flips involve flips of factors only, whereas signed flips involve flips of factors and of σ, provided the latter is defined.

Theorem 7.2. *Let P be a principal bundle with structure group $G = \mathrm{SO}(n)$ over a closed simply connected manifold of dimension 4. Two bundle reductions of P of type $[\mathrm{S}H^{\pm}]$ are conjugate under the action of the structure group if and only if they can be transformed into one another by a permutation of factors and by the following operations:*
– If there exists a signed stable factor or an O-factor of odd rank, any flip.
– Otherwise, unsigned flips and pairs of signed flips.

Proof. First, we consider bundle reductions of type $[\mathrm{S}H^{+}]$. Let H be a Howe subgroup of $\mathrm{O}(n)$ and let Q be a reduction of P to the subgroup $\mathrm{S}H^{+}$. Let $d \in \mathrm{SO}(n)$ such that $d(\mathrm{S}H^{+})d^{-1}$ is of standard form, i.e., equal to $\mathrm{S}K^{+}$ for some Howe subgroup K of $\mathrm{O}(n)$ of standard form. We have to determine how d acts on a connected component Q_0 of Q and derive from that how it acts on Q^H and on σ. We start with some preliminary considerations. The structure group of Q_0 is H_0. If $d(\mathrm{S}H^{+})d^{-1}$ is of standard form then so is its identity connected component dH_0d^{-1}. A similar argument as in the proof of Theorem 7.1 shows that then d can be decomposed as $d_1d_2d_3$, where d_1 is a pure permutation of the factors of H_0 (equivalently, those of H), $d_2 \in \mathrm{C}_{\mathrm{O}(n)}(H_0)$ and

$$d_3 = (a_1 \oplus \overset{k_1}{\cdots} \oplus a_1) \oplus \cdots \oplus (a_r \oplus \overset{k_r}{\cdots} \oplus a_r)\,,$$

with $a_i \in \mathrm{N}_{\mathrm{O}(\delta_i m_i)}\big((\mathrm{I}_{\mathbb{K}_i\, \overset{1}{m_i}})_0\big) \equiv \mathrm{N}_{\mathrm{O}(\delta_i m_i)}\big(\mathrm{I}_{\mathbb{K}_i\, \overset{1}{m_i}}\big)$. Since $d \in \mathrm{SO}(n)$,

$$(\det d_1)(\det d_2)(\det a_1)^{k_1} \cdots (\det a_r)^{k_r} = 1\,. \tag{56}$$

The only effect of d_2 is that it might compensate for a sign stemming from d_1 or the a_i. We claim that for this purpose it is sufficient that $d_2 \in \mathrm{C}_{\mathrm{O}(n)}(H)$. Indeed, since $\mathrm{C}_{\mathrm{O}(n)}(H_0)$ is a Howe subgroup, it contains an element of negative determinant if and only if it has an O-factor of odd multiplicity. Then the double centralizer $K = \mathrm{C}^2_{\mathrm{O}(n)}(H_0)$ contains an O-factor of odd rank. Since $H_0 \subseteq K \subseteq H$, K has the same dimension as H. It is therefore obtained from H by inverse field restriction of a factor O_2^k to the factor U_1^k or by merging a double factor $\mathrm{O}_1^{k_1} \times \mathrm{O}_1^{k_2}$ to the factor $\mathrm{O}_1^{k_1+k_2}$. In both cases, H has at least the same number of O-factors of odd rank as K, hence $\mathrm{C}_{\mathrm{O}(n)}(H)$ has an O-factor of odd multiplicity and therefore contains an element of negative determinant.

It follows that we may restrict attention to elements d of $\mathrm{SO}(n)$ for which $d_2 \in \mathrm{C}_{\mathrm{O}(n)}(H)$. For such elements, dHd^{-1} is of standard form. If, conversely, dHd^{-1} is of standard form then so is $d(\mathrm{S}H^+)d^{-1}$. This shows that the operations which apply to reductions of P to H_0 are given by those operations on reductions of $P^{\mathrm{O}(n)}$ to H_0 which can be implemented by $d \in \mathrm{SO}(n)$. We take the latter operations from the proof of Theorem 7.1 and determine what signs they require. Like there, let φ_i denote the automorphism of $\mathrm{I}_{\mathbb{K}_i}(m_i)$ induced by a_i. We check:

- The determinant of d_1 is given by the sign of the corresponding permutation of the standard basis in $\mathbb{R}^n$. If it is negative then H contains an O-factor of odd rank and odd multiplicity.
- The flips of signed factors of Q_0 produce a sign in (56): indeed, since the multiplicity is odd, it suffices to consider $\det a_i$. To induce a flip for the case of SO-factors of even rank and U-factors of odd rank, φ_i must be outer. Then Lemma 3.10 yields $\det a_i = -1$. The case of SO-factors of odd rank is degenerate in that the flip is trivial here. Nevertheless we may choose a_i so that $\det a_i = -1$ and assume that φ_i generates this (trivial) flip.
- The flips of unsigned factors of Q_0 do not produce a sign in (56): here either the multiplicity is even, in which case the assertion is obvious, or the factor is a U-factor of even rank, in which case the assertion follows from Lemma 3.10.

Now we can prove Theorem 7.2. If Q^H has an O-factor of odd rank, $\mathrm{C}_{\mathrm{O}(n)}(H)$ has an O-factor of odd multiplicity. Then $\mathrm{C}_{\mathrm{O}(n)}(H)$ contains an element of negative determinant so that we may choose the sign of d_2 appropriately to compensate for any sign produced by d_1 or the a_i in (56). It follows that in this case, any permutation and any flip can be applied to Q_0 and hence to Q^H and to σ. If Q^H does not have an O-factor of odd rank, $\det d_1 = 1$. In addition, $\mathrm{C}_{\mathrm{O}(n)}(H)$ does not have an O-factor of odd multiplicity, hence $\det d_2 = 1$. Hence, while permutations and unsigned flips of Q_0 can be applied arbitrarily, signed flips have to be paired. Since the properties of factors of Q_0 to be signed or unsigned and stable or unstable carry over to the corresponding factors of Q^H, signed/unsigned flips of U-factors of Q_0 translate into signed/unsigned flips of factors of Q^H, flips of signed unstable SO-factors translate into flips of σ and flips of unsigned SO-factors cancel. This proves the assertion for the case where Q^H does not possess a stable signed O-factor. If, on the other hand, Q^H has a stable signed factor then Q_0 has a stable signed factor. By pairing any signed flip with the flip of this factor we see that in effect any flip can be applied to Q_0 and hence to Q^H and to σ.

Finally, consider a bundle reduction Q of type $[\mathrm{S}H^-]$ of P. Let $d \in \mathrm{SO}(n)$ such that $d(\mathrm{S}H^-)d^{-1}$ is of standard form, i.e., equal to $\mathrm{S}K^-$ for some Howe subgroup K of $\mathrm{O}(n)$ of standard form. Let $a_{(n)}$ be the element of $\mathrm{O}(n)$ chosen to define $\mathrm{S}H^-$ and put $\tilde{d} = a_{(n)} d a_{(n)}^{-1}$. Let $\tilde{Q}$ be obtained by transforming Q by $a_{(n)}$ inside $Q^{\mathrm{O}(n)}$ and let $\tilde{P}$ be obtained by transforming P by $a_{(n)}$ inside $P^{\mathrm{O}(n)}$. Then $\tilde{Q}$ is a bundle reduction of type $[\mathrm{S}H^+]$ of $\tilde{P}$ and $d(\mathrm{S}H^-)d^{-1}$ is of standard form iff $\tilde{d}(\mathrm{S}H^+)\tilde{d}^{-1}$ is of standard form. Thus, any operation applicable to bundle reductions of type $[\mathrm{S}H^+]$ applies to $\tilde{Q}$ and hence to Q, by definition of the classifying data of Q. □

8 Summary: Classification of orbit types

We summarize the classification of orbit types obtained in this paper. For the sake of completeness, we include the results for $\mathrm{U}(n)$ and $\mathrm{SU}(n)$ derived in [34].

Let M be a closed, simply connected[7] manifold of dimension 4 and let P be a principal G-bundle over M , with G being a classical compact Lie group. Orbit types of the action of the group of

---

[7] The assumption that M be simply connected can be dropped in case $G = \mathrm{U}(n)$ or $\mathrm{SU}(n)$. For the other groups, it can be weakend to $H^1(M, \mathbb{Z}_2) = 0$, see Remark 4.1. In any case, the results hold as well for $\dim(M) = 2, 3$.

vertical automorphisms of P on the affine space of connections are 1:1 with isomorphism classes of holonomy-induced Howe subbundles of P, factorized by the action of the structure group.
In the following, r, m_i, k_i denote positive integers and w, p, e and c denote the Stiefel-Whitney, Pontryagin, Euler and Chern classes, respectively. Let $\tilde{p}(Q_i) = \sum_k (-1)^k p_k(Q_i)$ and $\tilde{c}(Q_i) = \sum_k (-1)^k c_k(Q_i)$ and let $\rho_2 : H^*(M, \mathbb{Z}) \to H^*(M, \mathbb{Z}_2)$ be induced by reduction mod 2.

Isomorphism classes of Howe subbundles of P

$\underline{G = \mathrm{U}(n):}$ The classes are labelled by

$$\big((m_1, k_1, c(Q_1)), \ldots, (m_r, k_r, c(Q_r))\big)\,, \tag{57}$$

where Q_i are principal $\mathrm{U}(m_i)$-bundles over M. These data are subject to the relations

$$n = \textstyle\sum_{i=1}^r k_i\, m_i\,, \tag{58}$$

$$c(P) = c(Q_1)^{k_1} \ldots c(Q_r)^{k_r}\,. \tag{59}$$

$\underline{G = \mathrm{SU}(n):}$ The classes are labelled by the data (57) and a characteristic class $\xi \in H^1(M, \mathbb{Z}_g)$ for $\mathbb{Z}_g$-bundles, where g is the greatest common divisor of the k_i. These data are subject to the relations (58), (59) and

$$\beta\xi = \textstyle\sum_{i=1}^r \frac{k_i}{g}\, c_1(Q_i)\,,$$

where $\beta : H^1(M, \mathbb{Z}_g) \to H^2(M, \mathbb{Z})$ is the connecting (Bockstein) homomorphism associated with the sequence of coefficient groups $0 \to \mathbb{Z} \to \mathbb{Z} \to \mathbb{Z}_g \to 0$.

$\underline{G = \mathrm{Sp}(n):}$ The classes are labeled by r-tuples

$$\big((\mathbb{K}_1, m_1, k_1, \alpha_1), \ldots, (\mathbb{K}_r, m_r, k_r, \alpha_r)\big)\,, \tag{60}$$

where $\mathbb{K}_i = \mathbb{R}$, $\mathbb{C}$ or $\mathbb{H}$ and α_i are characteristic classes of principal bundles Q_i over M with structure group $\mathrm{O}(m_i)$ for $\mathbb{K}_i = \mathbb{R}$, $\mathrm{U}(m_i)$ for $\mathbb{K}_i = \mathbb{C}$ and $\mathrm{Sp}(m_i)$ for $\mathbb{K}_i = \mathbb{H}$. In detail, $\alpha_i = \{w(Q_i), p(Q_i), [e](Q_i)\}$ for $\mathbb{K}_i = \mathbb{R}$ and $\alpha_i = c(Q_i)$ for $\mathbb{K}_i = \mathbb{C}$ or $\mathbb{H}$. Here $[e](Q_0)$ is defined as the equivalence class $\{\pm e(Q_{i0})\}$, where Q_{i0} is a reduction of Q_i to $\mathrm{SO}(m_i)$. These data are subject to the relation (58) and

$$c(P) = \textstyle\prod_{\mathbb{K}_i = \mathbb{R}}\, \tilde{p}(Q_i)^{2k_i} \cdot \prod_{\mathbb{K}_i = \mathbb{C}}\, c(Q_i)^{k_i}\, \tilde{c}(Q_i)^{k_i} \cdot \prod_{\mathbb{K}_i = \mathbb{H}}\, c(Q_i)^{k_i}\,. \tag{61}$$

$\underline{G = \mathrm{O}(n):}$ The classes are labeled by the data (60), subject to the relations

$$n = \textstyle\sum_{i=1}^r\, (\dim_{\mathbb{R}} \mathbb{K}_i)\, k_i\, m_i\,, \tag{62}$$

$$w(P) = \textstyle\prod_{\mathbb{K}_i = \mathbb{R}}\, w(Q_i)^{k_i} \cdot \prod_{\mathbb{K}_i = \mathbb{C},\mathbb{H}}\, \rho_2\,(c(Q_i))^{k_i}\,, \tag{63}$$

$$\tilde{p}(P) = \textstyle\prod_{\mathbb{K}_i = \mathbb{R}}\, \tilde{p}(Q_i)^{k_i} \cdot \prod_{\mathbb{K}_i = \mathbb{C},\mathbb{H}}\, c(Q_i)^{k_i}\, \tilde{c}(Q_i)^{k_i}\,, \tag{64}$$

$$[e](P) = \textstyle\prod_{\mathbb{K}_i = \mathbb{R}}\, [e](Q_i)^{k_i} \cdot \prod_{\mathbb{K}_i = \mathbb{C}}\, [c_{m_i}(Q_i)]^{k_i} \cdot \prod_{\mathbb{K}_i = \mathbb{H}}\, [c_{2m_i}(Q_i)]^{k_i}\,. \tag{65}$$

For $\mathbb{K}_i = \mathbb{R}$, the classes α_i are in addition subject to the relations valid for the characteristic classes of $\mathrm{O}(m_i)$-bundles, see Corollary 4.18 and the discussion after Equation (38).

$\underline{G = \mathrm{SO}(n):}$ The classes are labelled by those of the data (60) which satisfy
1. if there is i with $\mathbb{K}_i = \mathbb{R}$, $m_i = 2$ and k_i odd there is also $j \neq i$ with $\mathbb{K}_j = \mathbb{R}$ and k_j odd,
2. if there are $i \neq j$ with $\mathbb{K}_i = \mathbb{K}_j = \mathbb{R}$, $m_i = m_j = 1$ and k_i, k_j odd there is also $k \neq i, j$ with $\mathbb{K}_k = \mathbb{R}$ and k_k odd,

3. Conditions 1 and 2 with m and k exchanged.

Moreover, the following additional data occur. If there does not exist i with $\mathbb{K}_i = \mathbb{R}$ and m_i odd or k_i odd, or with $\mathbb{K}_i = \mathbb{C}$ and m_i, k_i odd, the Howe subgroup H of $\mathrm{O}(n)$ defined by the data (60) gives rise to two distinct conjugacy classes of Howe subgroups of $\mathrm{SO}(n)$, see Proposition 3.8. These two classes are distinguished by a sign $\pm$. If furthermore there exists i with $\mathbb{K}_i = \mathbb{R}$, k_i odd and $2e(Q_{i0}) \neq 0$ for some reduction Q_{i0} of Q_i to $\mathrm{SO}(m_i)$, the two reductions to the subgroup $\mathrm{SO}(n) \cap H$ of the principal bundle with structure group H defined by the data (60) are nonisomorphic. They are distinguished by a further sign $\sigma = \pm 1$. All of these data are subject to the relations (62)–(64) and

$$e(P) = \pm\sigma \cdot \prod\nolimits_{\mathbb{K}_i=\mathbb{R}} e^\circ(Q_i)^{k_i} \cdot \prod\nolimits_{\mathbb{K}_i=\mathbb{C}} c_{m_i}(Q_i)^{k_i} \cdot \prod\nolimits_{\mathbb{K}_i=\mathbb{H}} c_{2m_i}(Q_i)^{k_i}\,, \tag{66}$$

where $e^\circ(Q_i)$ are representatives for the equivalence classes $[e](Q_i)$. As for $\mathrm{O}(n)$, in case $\mathbb{K}_i = \mathbb{R}$, the characteristic classes are subject to the relations listed in Theorem 4.16.

Holonomy-induced Howe subbundles For $G = \mathrm{U}(n)$ or $\mathrm{SU}(n)$, all Howe subbundles are holonomy-induced. For $G = \mathrm{O}(n)$, $\mathrm{SO}(n)$, or $\mathrm{Sp}(n)$, a Howe subbundle is holonomy-induced if in the data (60) there is neither an i with $\mathbb{K}_i = \mathbb{R}$ and $m_i = 2$ nor an i and a $j \neq i$ with $\mathbb{K}_i = \mathbb{K}_j = \mathbb{R}$ and $m_i = m_j = 1$.

Factorization by the action of the structure group on bundle reductions The structure group acts on the data (60) and (57) by permuting the members $(\mathbb{K}_i, m_i, k_i, \alpha_i)$ and $(m_i, k_i, c(Q_i))$, respectively. For $G = \mathrm{Sp}(n)$, $\mathrm{O}(n)$ and $\mathrm{O}(n)$, the structure group acts in addition on the characteristic classes of the factors Q_i. For $G = \mathrm{Sp}(n)$ and $\mathrm{O}(n)$ this action is given by $c(Q_i) \mapsto \tilde{c}(Q_i)$. For $SO(n)$, we refer to Theorem 7.2.

9 Examples

To illustrate the above results, we will determine the Howe subbundles of principal bundles with structure groups $G = \mathrm{O}(4)$, $\mathrm{SO}(4)$, $\mathrm{Sp}(1)$ and $\mathrm{Sp}(2)$. We will start with $\mathrm{Sp}(1)$ and $\mathrm{Sp}(2)$, because there is only one equation to be studied here. To be definite, we restrict attention to the base manifolds $M = \mathrm{S}^2 \times \mathrm{S}^2$ and $\mathbb{C}\mathrm{P}^2$. Let us recall the ring structure of $H^*(M, \mathbb{Z})$ and $H^*(M, \mathbb{Z}_2)$. Let ζ denote the generator of $H^2(\mathrm{S}^2, \mathbb{Z})$ and let γ denote the generator of $H^2(\mathbb{C}\mathrm{P}^2, \mathbb{Z})$. Then

	$M = \mathrm{S}^2 \times \mathrm{S}^2$		$M = \mathbb{C}\mathrm{P}^2$	
	group	generated by	group	generated by
$H^2(M,\mathbb{Z})$	$\mathbb{Z} \oplus \mathbb{Z}$	$\zeta \times 1, 1 \times \zeta$	$\mathbb{Z}$	γ
$H^4(M,\mathbb{Z})$	$\mathbb{Z}$	$\zeta \times \zeta$	$\mathbb{Z}$	γ^2
$H^2(M,\mathbb{Z}_2)$	$\mathbb{Z}_2 \oplus \mathbb{Z}_2$	$\rho_2\zeta \times 1, 1 \times \rho_2\zeta$	$\mathbb{Z}_2$	$\rho_2\gamma$
$H^4(M,\mathbb{Z}_2)$	$\mathbb{Z}_2$	$\rho_2\zeta \times \rho_2\zeta$	$\mathbb{Z}_2$	$\rho_2\gamma^2$.

We use the following parameterization of characteristic classes of factors Q_i. For U- and Sp-factors,

$$c_1(Q_i) = x_i\,\gamma\,, \qquad c_2(Q_i) = z_i\,\gamma^2$$

in case $M = \mathbb{C}\mathrm{P}^2$ and

$$c_1(Q_i) = x_i\,\zeta \times 1 + y_i\,1 \times \zeta\,, \qquad c_2(Q_i) = z_i\,\zeta \times \zeta$$

in case $M = \mathrm{S}^2 \times \mathrm{S}^2$, where x_i, y_i, z_i are integers. The only O-factors we will come about have structure group $\mathrm{O}(2)$ or $\mathrm{O}(3)$. In case of an $\mathrm{O}(2)$-factor, $w_2(Q_i)$ and $p_1(Q_i)$ can be expressed in terms of $[e](Q_i)$, see Theorem 4.16 and Corollary 4.18. Writing, respectively,

$$[e](Q_i) = f_i\,[\gamma]\,, \qquad [e](Q_i) = [f_i\,\zeta \times 1 + g_i\,1 \times \zeta]$$

with integers $f_i, g_i \in \mathbb{Z}$, we obtain

$$w_2(Q_i) = f_i\,\rho_2\gamma\,, \qquad p_1(Q_i) = f_i^2\,\gamma^2$$

for $M = \mathbb{C}\mathrm{P}^2$ and

$$w_2(Q_i) = f_i\,\rho_2\zeta \times 1 + g_i\,1 \times \rho_2\zeta\,, \qquad p_1(Q_i) = 2 f_i g_i\,\zeta \times \zeta$$

for $M = \mathrm{S}^2 \times \mathrm{S}^2$. The integers f_i, g_i are determined uniquely if we require $f_i \geq 0$ or $(f_i, g_i) \geq (0, 0)$ (lexicographic ordering), respectively. In case of an $\mathrm{O}(3)$-factor, we write

$$w_2(Q_i) = s_i\,\rho_2\gamma\,, \qquad p_1(Q_i) = a_i\,\gamma^2$$

in case $M = \mathbb{C}\mathrm{P}^2$ and

$$w_2(Q_i) = s_i\,\rho_2\zeta \times 1 + t_i\,1 \times \rho_2\zeta\,, \qquad p_1(Q_i) = a_i\,\zeta \times \zeta$$

in case $M = \mathrm{S}^2 \times \mathrm{S}^2$, where $s_i, t_i, a_i \in \mathbb{Z}$. The parameters s_i, t_i are uniquely determined if we require $s_i, t_i = 0, 1$.
Finally, let us note the following. With the exception of $\mathrm{SO}(2)$, the center of $\mathrm{O}(n)$, $\mathrm{SO}(n)$ or $\mathrm{Sp}(n)$ (corresponding to the Howe subgroup O_1^n) is finite. Due to the assumption that M be simply connected, reductions to the center are therefore necessarily trivial and hence occur exactly when P is trivial. They will not be mentioned below.

9.1 Sp(1) and Sp(2)

The reduction equation is given by (53). We parameterize the characteristic class of P by an integer Z, i.e., $c_2(P) = Z\gamma^2$ for $\mathbb{C}\mathrm{P}^2$ and $c_2(P) = Z\zeta \times \zeta$ for $\mathrm{S}^2 \times \mathrm{S}^2$. According to Theorem 7.1, reductions have to be identified iff they can be transformed into one another by flips of U-factors and permutations. To factorize by flips we require $x_i \geq 0$ or $(x_i, y_i) \geq (0, 0)$ (lexicographic ordering), respectively, for any U-factor. To factorize by permutations we fix an order of the factors H_i for each conjugacy class of Howe subgroups H and require that the parameters of identical factors (same field, rank and multiplicity) increase w.r.t. lexicographic ordering. In detail, for $i < j$ we require

- $f_i \leq f_j$ or $(f_i, g_i) \leq (f_j, g_j)$, respectively, whenever $H_i = H_j = \mathrm{O}_2^k$,
- $(a_i, s_i) \leq (a_j, s_j)$ or $(a_i, s_i, t_i) \leq (a_j, s_j, t_j)$, respectively, whenever $H_i = H_j = \mathrm{O}_3^k$; here the restriction on s_i, s_j, t_i, t_j to take the values 0 and 1 only is understood,
- $(x_i, z_i) \leq (x_j, z_j)$ or $(x_i, y_i, z_i) \leq (x_j, y_j, z_j)$, respectively, whenever $H_i = H_j = \mathrm{U}_m^k$,
- $z_i \leq z_j$ whenever $H_i = H_j = \mathrm{Sp}_m^k$.

Now consider $G = \mathrm{Sp}(1)$. The only nontrivial Howe subgroup is U_1^1. It is holonomy-induced. Parameters are $x \in \mathbb{Z}$ for $M = \mathbb{C}\mathrm{P}^2$ and $x, y \in \mathbb{Z}$ for $M = \mathrm{S}^2 \times \mathrm{S}^2$. The reduction equation (53) reads $c(P) = c(Q)\tilde{c}(Q)$. In terms of parameters, this amounts to

$$Z = -x^2 \qquad \text{or} \qquad Z = -2xy\,,$$

respectively. In case $M = \mathbb{C}\mathrm{P}^2$, a solution x exists iff $-Z$ is a square. Due to $0 \leq x$, the solution is unique. In case $M = \mathrm{S}^2 \times \mathrm{S}^2$, solutions exist iff Z is even. Under the condition $(0,0) \leq (x,y)$ the solutions are parameterized as follows. If $Z \neq 0$, $x = \varepsilon_1, \cdots, \varepsilon_s$, where ε_l are the (nonnegative) divisors of $\frac{|Z|}{2}$. If $Z = 0$, one has $x = y = 0$ and the two families of solutions $x > 0$, $y = 0$ and $x = 0$, $y > 0$. Thus, over $\mathbb{C}\mathrm{P}^2$ the number of orbit types with stabilizer isomorphic to U(1) can be 0 or 1, whereas over $\mathrm{S}^2 \times \mathrm{S}^2$ it can be any nonnegative integer or countably infinite.

Remark 9.1. By virtue of the isomorphism $\varphi_{\mathbb{H},\mathbb{C}} : \mathrm{Sp}(1) \to \mathrm{SU}(2)$, U_1^1 is mapped to the subgroup of diagonal matrices in SU(2). The above result is consistent with what is known about reductions of SU(2)-bundles to this subgroup.

Now consider $G = \mathrm{Sp}(2)$. Equations will be given in terms of parameters only. First, we discuss holonomy-induced Howe subbundles, see Figure 5.

U_1^2: In case $M = \mathbb{C}\mathrm{P}^2$, $Z = -2x^2$; in case $M = \mathrm{S}^2 \times \mathrm{S}^2$, $Z = -4xy$. The discussion is analogous to the case of the Howe subgroup U_1^1 of Sp(1).

$\mathrm{O}_1^1 \times \mathrm{U}_1^1$: In case $M = \mathbb{C}\mathrm{P}^2$, $Z = -x^2$. In case $M = \mathrm{S}^2 \times \mathrm{S}^2$, $Z = -2xy$. The parameters x and y belong to the U-factor. In both cases, the discussion is analogous to the case of the Howe subgroup U_1^1 of Sp(1).

$\mathrm{O}_1^1 \times \mathrm{Sp}_1^1$: In both cases, the equation is $Z = z$, where z belongs to the Sp-factors. Hence, there exists a unique reduction for any P.

Sp_1^2: In both cases, the equation is $Z = 2z$. There exists a reduction iff Z is even and the reduction is unique.

$\mathrm{Sp}_1^1 \times \mathrm{Sp}_1^1$: In both cases, the equation is $Z = z_1 + z_2$. Reductions exist for any P. They can be parameterized by $z_1 \in \mathbb{Z}$.

$\mathrm{U}_1^1 \times \mathrm{U}_1^1$: In case $M = \mathbb{C}\mathrm{P}^2$, $Z = -(x_1^2 + x_2^2)$. There is no reduction for $Z > 0$ and a unique reduction $x_1 = x_2 = 0$ for $Z = 0$. For $Z < 0$, there is the classical result that for given integer n the number of solutions of the equation $a^2 + b^2 = n$ satisfying $a \geq 0$ and $b > 0$ is given by $N(n) = \sum_{d|n} \chi(d)$, where the sum runs over the divisors of n and χ denotes the nontrivial Dirichlet character modulo 4, defined by $\chi(d) = 1$ if $d = 1 \mod 4$, $\chi(d) = -1$ if $d = 3 \mod 4$ and $\chi(d) = 0$ if d is even [24]. First, it follows that reductions exist if and only if those of the prime factors p of $-Z$ which satisfy $p = 3 \mod 4$ appear with even powers. That this condition is sufficient for the existence of reductions can also be deduced directly from Fermat's theorem on sums of two squares and the Brahmagupta-Fibonacci identity. Second, it follows that the number of reductions, i.e., solutions satisfying $0 \leq x_1 \leq x_2$, is $\frac{N(-Z)+1}{2}$ if $-Z$ is a square or double a square and $\frac{N(-Z)}{2}$ otherwise. For example, for small n we have $N(-Z) = 0$ for $-Z = 3, 6$, $N(-Z) = 1$ for $-Z = 1, 2, 4, 8, 9$ and $N(-Z) = 2$ for $-Z = 5, 10$; yet the reduction is unique also in the last two cases. On the other hand, it is easy to see that both $N(-Z)$ and the number of reductions can take any nonnegative integer value.
In case $M = \mathrm{S}^2 \times \mathrm{S}^2$, $Z = -2(x_1 y_1 + x_2 y_2)$. Reductions exist iff Z is even. They may be enumerated by choosing $(x_1, y_1) \geq (0,0)$ arbitrarily and letting (x_2, y_2) run through the solutions of $x_2 y_2 = -\frac{Z}{2} - x_1 y_1$ which satisfy $(x_2, y_2) \geq (x_1, y_1)$ (for certain values of (x_1, y_1) there may be no such solutions).

$\mathrm{U}_1^1 \times \mathrm{Sp}_1^1$: In case $M = \mathbb{C}\mathrm{P}^2$, $Z = z - x^2$. In case $M = \mathrm{S}^2 \times \mathrm{S}^2$, $Z = z - 2xy$. Here x and y belong to the U-factor and z belongs to the Sp-factor. Reductions exist for all P. They are parameterized by $x \geq 0$ in case $M = \mathbb{C}\mathrm{P}^2$ and $(x,y) \geq 0$ in case $M = \mathrm{S}^2 \times \mathrm{S}^2$.

U_2^1: In case $M = \mathbb{C}\mathrm{P}^2$, $Z = 2z - x^2$. Reductions always exist. They are parameterized by $x = 2l$ if Z is even and $x = 2l+1$ if Z is odd, where $l = 0, 1, 2, \ldots$. In case $M = \mathrm{S}^2 \times \mathrm{S}^2$, $Z = 2(z - xy)$. Reductions exist iff Z is even. They are parameterized by $(x, y) \geq (0, 0)$ then.

Out of the Howe subgroups whose reductions are not holonomy-induced, there is only one which does not consist exclusively of O(1)-factors:

O_2^1: Here the abstract reduction equation is $c(P) = \tilde{p}(Q)^2$. Using the relation $p_1(Q) = [e](Q)^2$, see Theorem 4.16 and Corollary 4.18, we obtain $c_2(P) = -2[e](Q)^2$. In case $M = \mathbb{C}\mathrm{P}^2$, this equation becomes $Z = -2f^2$ with $f \geq 0$. In case $M = \mathrm{S}^2 \times \mathrm{S}^2$, $Z = -4fg$ with $(f, g) \geq (0, 0)$. The discussion of these equations is analogous to the case of the Howe subgroup U_1^1 of Sp(1).

9.2 O(4)

We parameterize the characteristic classes of P by

$$w_2(P) = S\,\rho_2\gamma\,, \qquad p_1(P) = A\gamma^2\,, \qquad [e](P) = F[\gamma^2] \tag{67}$$

for $M = \mathbb{C}\mathrm{P}^2$ and by

$$w_2(P) = S\,\rho_2\zeta \times 1 + T\,1 \times \rho_2\zeta\,, \qquad p_1(P) = A\zeta \times \zeta\,, \qquad [e](P) = F[\zeta \times \zeta] \tag{68}$$

for $M = \mathrm{S}^2 \times \mathrm{S}^2$, where A, F, S, T are integers. S, T, F are determined uniquely under the conditions $S, T = 0, 1$ and $F \geq 0$. These conditions will be assumed to hold below. The characteristic classes of P are subject to a single relation which arises from the relation $w_4(P) = \rho_2[e](P)$ of Theorem 4.16 by replacing $w_4(P)$ by means of the fundamental relation, see (38). Using (87) we find that in terms of the parameters, this relation reads, respectively,

$$A - S^2 - 2F = 0 \mod 4\,, \qquad A - 2ST - 2F = 0 \mod 4\,. \tag{69}$$

In particular, in case $M = \mathrm{S}^2 \times \mathrm{S}^2$, A must be even. To obtain unique representatives under the action of the structure group, see Theorem 7.1, the same conditions on the parameters as for the structure groups Sp(1) and Sp(2) above have to be imposed. The reduction equations are given by (46), (47), (49). We discuss them in terms of the parameters, starting with the holonomy-induced reductions. The equation for $w_4(P)$ needs not be considered, because it is automatically satisfied if the equation for $[e](P)$ holds.

U_1^2: In case $M = \mathbb{C}\mathrm{P}^2$, $S = 0 \mod 2$, $A = 2x^2$, $F = \pm x^2$. Reductions exist iff $S = 0 \mod 2$, $A = 2F$ and F is a square. Under the condition $x \geq 0$, they are unique. In case $M = \mathrm{S}^2 \times \mathrm{S}^2$, $S = T = 0$, $A = 4xy$ and $F = \pm 2xy$. Reductions exist iff $S = T = 0$ and $A = \pm 2F$. Under the condition $(x, y) \geq (0, 0)$, they are parameterized as follows. If $F \neq 0$, $(x, y) = (\varepsilon, \pm\frac{F}{2\varepsilon})$, where ε runs through the (nonnegative) divisors of F. If $F = 0$, one has $x = y = 0$ and the two families of solutions $x > 0$, $y = 0$ and $x = 0$, $y > 0$.

$\mathrm{O}_1^2 \times \mathrm{U}_1^1$: In case $M = \mathbb{C}\mathrm{P}^2$, $S = x \mod 2$, $A = x^2$ and $F = 0$ where the parameter x refers to the U-factor. A reduction exists iff $F = 0$ and A is a square. Due to $x \geq 0$, it is unique. Since the mod 4 reduction of a square is 0 if the square is even and 1 if the square is odd, the equation for S is automatically satisfied due to (69).
In case $M = \mathrm{S}^2 \times \mathrm{S}^2$, $S = x \mod 2$, $T = y \mod 2$, $A = 2xy$ and $F = 0$. Reductions exist iff $F = 0$ and ($S = 1$ or $T = 1$ or A is divisible by 8). If $A = 0$, (69) implies $S = 0$ or $T = 0$. Accordingly, reductions are parameterized by $x = 0$, $y = 2l + T$ or $y = 0$, $x = 2l + S$, $l \geq 0$. If $A \neq 0$, reductions are parameterized as follows. If $S = T = 0$, A must be divisible by 8 and

$x = 2\varepsilon_1, \ldots, 2\varepsilon_s$, where ε_i are the divisors of $\frac{A}{8}$. If $S = 1$, $T = 0$, x runs through the odd divisors of $\frac{A}{2}$. Similarly for y in case $S = 0$, $T = 1$. If $S = T = 1$, x runs through the divisors of $\frac{A}{2}$.

$\mathrm{O}_1^1 \times \mathrm{O}_3^1$: In case $M = \mathbb{C}\mathrm{P}^2$, $S = s \mod 2$, $A = a$, $F = 0$ where the parameters refer to the O(3)-factor. Together with (69), these equations imply $a - s^2 = 0 \mod 4$, which is the fundamental relation for the O(3)-bundle parameterized by a and s. Hence, reductions exist iff $F = 0$ and are unique then. In case $M = \mathrm{S}^2 \times \mathrm{S}^2$, the above equations hold, together with $T = t \mod 2$. The result is analogous.

Sp_1^1: Here, $A = 2z$, $F = \pm z$, $S = 0$ or $S = T = 0$, respectively. Reductions exist iff $S = 0$ or $S = T = 0$, respectively, and $A = \pm 2F$. They are unique.

U_2^1: In case $M = \mathbb{C}\mathrm{P}^2$, $S = x \mod 2$, $A = x^2 - 2z$, $F = \pm z$. Reductions exist iff $A - 2F$ or $A + 2F$ is a square. They are unique in each of these two cases. (That means in particular, if both $A - 2F$ or $A + 2F$ are squares and $F \neq 0$ there exist exactly 2 solutions.) By the same argument as in the discussion of the Howe subgroup $\mathrm{O}_1^2 \times \mathrm{U}_1^1$, (69) implies that the equation for S is automatically satisfied.
In case $M = \mathrm{S}^2 \times \mathrm{S}^2$, $S = x \mod 2$, $T = y \mod 2$, $A = 2(xy - z)$, $F = \pm z$. The last and the third equation yield $xy = \frac{A}{2} \pm F$. Hence, in case $F \neq 0$, potentially there are two families of reductions, corresponding to the two signs. For each of these families, the discussion is analogus to the case of reductions to the Howe subgroup $\mathrm{O}_1^2 \times \mathrm{U}_1^1$ above, with A replaced by $A \pm 2F$.

$\mathrm{U}_1^1 \times \mathrm{U}_1^1$: In case $M = \mathbb{C}\mathrm{P}^2$,

$$S = (x_1 + x_2) \mod 2\,, \qquad A = x_1^2 + x_2^2\,, \qquad F = x_1 x_2\,. \tag{70}$$

We have omitted the equation $F = -x_1 x_2$, because it does not have a solution due to the requirement $0 \leq x_1 \leq x_2$. Define $x_\pm := x_2 \pm x_1$. Then (70) translates into

$$S = x_+ \mod 2\,, \qquad A + 2F = x_+^2\,, \qquad A - 2F = x_-^2 \tag{71}$$

and the requirement $0 \leq x_1 \leq x_2$ translates into the requirement $0 \leq x_- \leq x_+$. Systems (70) and (71) are equivalent: the only thing to be checked is that if $x_\pm$ is a solution of (71) then $x_1 = \frac{x_+ - x_-}{2}$ and $x_2 = \frac{x_+ + x_-}{2}$ are integers. Since x_+^2 and x_-^2 differ by $4F$, this is obvious. Thus, a reduction exists iff $A = S \mod 2$ and $A \pm 2F$ are both squares. If it exists, it is unique.
In case $M = \mathrm{S}^2 \times \mathrm{S}^2$,

$$\begin{aligned} S &= (x_1 + x_2) \mod 2\,, & T &= (y_1 + y_2) \mod 2\,, \\ A &= 2(x_1 y_1 + x_2 y_2)\,, & F &= \pm(x_1 y_2 + x_2 y_1) \end{aligned} \tag{72}$$

Define $x_\pm := x_2 \pm x_1$ and $y_\pm := y_2 \pm y_1$. Then (72) translates into

$$\begin{aligned} S = x_+ = x_- \mod 2\,, & \qquad\qquad & T = y_+ = y_- \mod 2\,, \\ \tfrac{A}{2} + F = x_+ y_+\,, \quad \tfrac{A}{2} - F = x_- y_- & \qquad \text{or} \qquad & \tfrac{A}{2} + F = x_- y_-\,, \quad \tfrac{A}{2} - F = x_+ y_+ \end{aligned} \tag{73}$$

(recall that A is now even due to the relation (69); moreover we have added the obvious equations $S = x_- \mod 2$ and $T = y_- \mod 2$). The requirement $(0,0) \leq (x_1, y_1) \leq (x_2, y_2)$ translates into the requirement $(0,0) \leq (x_-, y_-) \leq (x_+, y_+)$. It is straightforward to check that systems (72) and (73) are equivalent. We discuss the solutions of the latter system.

– If $S = T = 1$, the relation (69) implies that $\frac{A}{2} \pm F$ is odd. Hence, reductions always exist. They are parameterized by arbitrary combinations of decompositions $\frac{A}{2} + F = a_+ b_+$

and $\frac{A}{2} - F = a_- b_-$ with $(a_\pm, b_\pm) \geq (0,0)$. Then $(x_+, y_+) = \max\{(a_+, b_+), (a_-, b_-)\}$ and $(x_-, y_-) = \min\{(a_+, b_+), (a_-, b_-)\}$

– If $S = 1$, $T = 0$, the relation (69) implies that $\frac{A}{2} \pm F$ is even. Therefore, again, reductions always exist. They are parameterized in the same way as in the case $S = T = 1$, with the additional condition that $a_\pm$ has to be odd.

– The case $S = 0$ and $T = 1$ is similar. Instead of $a_\pm$, $b_\pm$ has to be odd.

– If $S = T = 0$, the relation (69) yields that $\frac{A}{2} \pm F$ is even. For reductions to exist, however, both $\frac{A}{2} + F$ and $\frac{A}{2} - F$ have to be divisible by 4. In this case, reductions are parameterized by arbitrary combinations of decompositions $\frac{A+2F}{8} = a_+ b_+$ and $\frac{A-2F}{8} = a_- b_-$ with $(a_\pm, b_\pm) \geq (0,0)$. Then $(x_+, y_+) = \max\{(2a_+, 2b_+), (2a_-, 2b_-)\}$ and $(x_-, y_-) = \min\{(2a_+, 2b_+), (2a_-, 2b_-)\}$

Next, we discuss the reductions which are not holonomy-induced.

O_2^2: In case $M = \mathbb{C}\mathrm{P}^2$, $S = 0$, $A = 2f^2$, $F = \pm f^2$; in case $M = \mathrm{S}^2 \times \mathrm{S}^2$, $S = T = 0$, $A = 4fg$, $F = \pm 2fg$. The discussion is analogous to that for the Howe subgroup U_2^1, with x, y replaced by f, g[8]. The present requirement $f \geq 0$ or $(f, g) \geq (0,0)$, respectively, which ensures that the parameters are uniquely determined corresponds to the requirement $x \geq 0$ or $(x, y) \geq (0,0)$, respectively, which singles out unique representatives for the classes of bundle reductions under the action of the structure group there.

$\mathrm{O}_2^1 \times \mathrm{O}_2^1$: In case $M = \mathbb{C}\mathrm{P}^2$, $S = (f_1 + f_2) \mod 2$, $A = f_1^2 + f_2^2$, $F = \pm f_1 f_2$; in case $M = \mathrm{S}^2 \times \mathrm{S}^2$, $S = (f_1 + f_2) \mod 2$, $T = (g_1 + g_2) \mod 2$, $A = 2(f_1 g_1 + f_2 g_2)$, $F = \pm(f_1 g_2 + f_2 g_1)$. The discussion is analogous to that for the Howe subgroup $\mathrm{U}_1^1 \times \mathrm{U}_1^1$, with x_i, y_i replaced by f_i, g_i.

$\mathrm{O}_2^1 \times \mathrm{U}_1^1$: In case $M = \mathbb{C}\mathrm{P}^2$, $S = (f + x) \mod 2$, $A = f^2 + x^2$, $F = \pm fx$; in case $M = \mathrm{S}^2 \times \mathrm{S}^2$, $S = (f + x) \mod 2$, $T = (g + y) \mod 2$, $A = 2(fg + xy)$, $F = \pm(fy + xg)$. Variables f, g refer to the O-factor, variables x, y to the U-factor. The discussion is analogous to that for the Howe subgroup $\mathrm{U}_1^1 \times \mathrm{U}_1^1$, with x_1, y_1 replaced by f, g and x_2, y_2 replaced by x, y; the only difference is that here permutations do not appear, hence the variables f, g and x, y are independent from one another.

$\mathrm{O}_2^1 \times \mathrm{O}_1^2$: In case $M = \mathbb{C}\mathrm{P}^2$, $S = f \mod 2$, $A = f^2$, $F = 0$; in case $M = \mathrm{S}^2 \times \mathrm{S}^2$, $S = f \mod 2$, $T = g \mod 2$, $A = 2fg$, $F = 0$. The parameters refer to the last factor. The discussion is analogous to that for the Howe subgroup $\mathrm{O}_1^2 \times \mathrm{U}_1^1$, with x, y replaced by f, g.

For $\mathrm{O}_1^1 \times \mathrm{O}_1^1 \times \mathrm{U}_1^1$ and $\mathrm{O}_1^1 \times \mathrm{O}_1^1 \times \mathrm{O}_2^1$, see $\mathrm{O}_1^2 \times \mathrm{U}_1^1$.

9.3 SO(4)

The characteristic classes $w_2(P)$ and $p_1(P)$ are parameterized as in (67) and (68). The Euler class will be parameterized by $e(P) = F\gamma^2$ in case $M = \mathbb{C}\mathrm{P}^2$ and by $e(P) = F\gamma \times \gamma$ in case $M = \mathrm{S}^2 \times \mathrm{S}^2$. Here F can take any integer value. Again, the characteristic classes of P are subject to the single relation (69). The reduction equations are given by (46), (47) and (50). In comparison with the case of structure group O(4), the corresponding equations for the parameters A, S and T are the same, whereas the equation for F is modified as follows:

– F can take any integer value,

– The sign is either positive or negative, according to whether the Howe subgroup $\mathrm{S}H^+$ or $\mathrm{S}H^-$ is considered. Thus, solutions for positive sign belong to $\mathrm{S}H^+$ and solutions for negative sign belong to $\mathrm{S}H^-$. If $\mathrm{S}H^+$ and $\mathrm{S}H^-$ are conjugate in $\mathrm{SO}(4)$, only the equation with the positive sign has to be taken into account.

---

[8] In fact, $[e](Q)$ here and $c_1(Q)$ there are related via the extension of the structure group to O(2). This holds similarly for the other reductions which are not holonomy-induced.

- The factor σ is added.

First, we discuss holonomy-induced Howe subbundles. Here, $\sigma = 1$, because if $G = \mathrm{SO}(4)$ then $\sigma = -1$ requires an $\mathrm{O}(2)$-factor. The system of equations is therefore the same as for the corresponding subgroup of $\mathrm{O}(4)$. We comment on how the discussion has to be modified and we derive the conditions on the parameters of the reductions which have to be imposed in order to obtain unique representatives w.r.t. the action of the structure group $\mathrm{SO}(4)$. Generally, $F \in \mathbb{Z}$ and solutions for positive sign belong to $\mathrm{S}H^+$ and solutions for negative sign belong to $\mathrm{S}H^-$.

$\mathrm{U}_1^{2\,\pm}$: In case $M = \mathbb{C}\mathrm{P}^2$ the condition $\langle A = 2F$ and F is a square$\rangle$ has to be replaced by $\langle A = \pm 2F$ and $\pm F$ is a square$\rangle$. In particular, in the case of $\mathrm{U}_1^{2\,+}$, necessarily $F \geq 0$ whereas in the case of $\mathrm{U}_1^{2\,-}$, $F \leq 0$. Since the factor $\mathrm{U}_1^{2\,\pm}$ is unsigned, any flip is allowed. Hence, we have to require $x \geq 0$ in case $M = \mathbb{C}\mathrm{P}^2$ and $(x, y) \geq (0, 0)$ in case $M = \mathrm{S}^2 \times \mathrm{S}^2$. This is the same condition as in the case of structure group $\mathrm{O}(4)$.

$\mathrm{U}_1^1 \times \mathrm{U}_1^1$: Only the positive sign has to be discussed. In case $M = \mathbb{C}\mathrm{P}^2$ this was so before for another reason. In case $M = \mathrm{S}^2 \times \mathrm{S}^2$ this amounts to setting $(x_+, y_+) = (a_+, b_+)$ and $(x_-, y_-) = (a_-, b_-)$. Since the factors of $\mathrm{U}_1^1 \times \mathrm{U}_1^1$ are both signed, flips can be applied in pairs only. We thus have to require $(x_1, x_2) \geq (0, 0)$ in case $M = \mathbb{C}\mathrm{P}^2$ and $((x_1, y_1), (x_2, y_2)) \geq ((0, 0), (0, 0))$ (lexicographic ordering) in case $M = \mathrm{S}^2 \times \mathrm{S}^2$. We leave it to the reader to translate these conditions to $(x_\pm, y_\pm)$.

$\mathrm{Sp}_1^{1\,\pm}$: The discussion of the reduction equations is modified in the standard way. The action of the structure group $\mathrm{SO}(4)$ is trivial.

$\mathrm{U}_2^{1\,\pm}$: Again, the discussion of the reduction equations is modified in the standard way. Since the factor U_2^1 is unsigned, it can be flipped. This leads to the same conditions on the parameters as in the case of $\mathrm{O}(4)$.

Next, we discuss Howe subbundles which are not holonomy-induced, omitting Howe subgroups consisting enirely of $\mathrm{O}(1)$-factors. To define the invariant σ, as unique representatives for the classes in $\mathrm{P}H^2(M, \mathbb{Z})$ we choose the elements of $H^2(M, \mathbb{Z})$ with nonnegative (w.r.t. lexicographic ordering) coefficients w.r.t. the generators γ and $\gamma \times 1$, $1 \times \gamma$, respectively. For an integer x, define $\mathrm{sgn}(x) = 1$ if $x \geq 0$ and $\mathrm{sgn}(x) = -1$ otherwise. For integers x, y define $\mathrm{sgn}(x, y) = 1$ if $(x, y) \geq (0, 0)$ and $\mathrm{sgn}(x, y) = -1$ otherwise.

$\mathrm{O}_2^{2\,\pm}$: Since the multiplicity is even, $\sigma = 1$. Therefore, the reduction equations are the same as for $\mathrm{O}(4)$ and the discussion is modified in the standard way. The action of the structure group $\mathrm{SO}(4)$ is trivial.

$\mathrm{O}_2^1 \times \mathrm{O}_2^1$: In the equation for F, σ can be omitted if one allows f_i and (f_i, g_i), respectively, to take arbitrary values. This amounts to seeking reductions to the identity connected component, which coincides with the Howe subgroup $\mathrm{U}_1^1 \times \mathrm{U}_1^1$ discussed above. For any reduction (x_1, y_1), (x_2, y_2) found there, we define:

- in case $M = \mathbb{C}\mathrm{P}^2$, $\sigma = \mathrm{sgn}(x_1)\,\mathrm{sgn}(x_2)$ and $f_i = \mathrm{sgn}(x_i) x_i$, $i = 1, 2$,
- in case $M = \mathrm{S}^2 \times \mathrm{S}^2$, $\sigma = \mathrm{sgn}(x_1, y_1)\,\mathrm{sgn}(x_2, y_2)$ and $(f_i, g_i) = \mathrm{sgn}(x_i, y_i)(x_i, y_i)$, $i = 1, 2$.

This way, due to the conditions imposed on x_i and (x_i, y_i), respectively, each reduction to $\mathrm{O}_2^1 \times \mathrm{O}_2^1$ arises from exactly one reduction to $\mathrm{U}_1^1 \times \mathrm{U}_1^1$. For the same reason, the action of the structure group $\mathrm{SO}(4)$ is already factored out. For completeness, let us just state what this action amounts to. As operations there occur the interchange of factors and the flip of σ. Since both factors of $\mathrm{O}_2^1 \times \mathrm{O}_2^1$ are signed, flips have to be applied in pairs which means that σ remains unchanged unless one of the factors is stable and the other one is not. That is, $f_1 = 0$ and $f_2 \neq 0$ or $(f_1, g_1) = (0, 0)$ and $(f_2, g_2) \neq (0, 0)$, respectively, or vice versa.

Acknowledgements

The authors are very much indebted to M. Čadek for helpful hints concerning the classification of real vector bundles. They are also grateful to J. Huebschmann for discussing with them several tools from algebraic topology used in the paper.

A. Hertsch acknowledges financial support by the International Max Planck Research School of the Max Planck Institute for Mathematics in the Sciences (MPI-MIS) at Leipzig.

A Universal bundles

We collect some basic facts about universal bundles and classifying spaces. For details, see e.g. [41, §19] or [23].

For every Lie group G and every positive integer n there exists a principal G-bundle $(\mathrm{E}\,G)_n \to (\mathrm{B}\,G)_n$ such that for every connected $(n-1)$-dimensional CW-complex X, the assignment of the pull-back bundle $f^*(\mathrm{E}\,G)_n$ to a map $f : X \to (\mathrm{B}\,G)_n$ induces a bijection from the set of homotopy classes of maps $X \to (\mathrm{B}\,G)_n$ onto the set of isomorphism classes of principal G-bundles over X. The bundle $(\mathrm{E}\,G)_n$ is referred to as an n-universal bundle for G, $(\mathrm{B}\,G)_n$ is referred to as an n-classifying space of G and the map $f : X \to (\mathrm{B}\,G)_n$ associated with a principal G-bundle P over X is referred to as an n-classifying map of P. Given a principal G-bundle $E \to B$, this bundle is n-universal for G if and only if B is path connected and

$$\pi_i(E) = 0\,, \qquad 1 \leq i < n\,. \tag{74}$$

For the groups considered in this paper, n-universal bundles are constructed as follows. For $\mathbb{K} = \mathbb{R}$, $\mathbb{C}$, $\mathbb{H}$ and positive integers $k < l$ let $V^{\mathbb{K}}_{l,k}$ denote the Stiefel manifold of orthonormal k-frames in $\mathbb{K}^l$ and let $G^{\mathbb{K}}_{l,k}$ denote the Grassmann manifold of k-dimensional subspaces in $\mathbb{K}^l$. By mapping an orthonormal frame to the subspace it spans one obtains a principal bundle $V^{\mathbb{K}}_{l,k} \to G^{\mathbb{K}}_{l,k}$ with structure group $\mathrm{I}_{\mathbb{K}}(k)$ (see Section 3 for this notation), known as the $\mathbb{K}$-Stiefel bundle of k-frames in dimension l. Since $\pi_i(V^{\mathbb{K}}_{l,k}) = 0$ for $i < (\dim_{\mathbb{R}} \mathbb{K})\,(l - k + 1) - 1$, the Stiefel bundle in dimension $l = m + k - 1$ is n-universal for $\mathrm{I}_{\mathbb{K}}(k)$, where $n = (\dim_{\mathbb{R}} \mathbb{K})\, m - 1$.

The n-universal bundles of a Lie group G can be arranged such that there exist embeddings $(\mathrm{E}\,G)_{n_i} \hookrightarrow (\mathrm{E}\,G)_{n_{i+1}}$ for an increasing sequence $n_1 < n_2 < \dots$. The inductive limit of such a sequence provides a universal bundle $\mathrm{E}\,G \to \mathrm{B}\,G$ for the group G. A general construction of the universal bundle as the infinite join of G is given in [31], see also [23]. Since $\pi_i(\mathrm{E}\,G) = 0$ for all i, the long exact homotopy sequence of the universal bundle implies

$$\pi_i(G) = \pi_{i+1}(\mathrm{B}\,G)\,, \qquad i = 0, 1, 2, \dots\ . \tag{75}$$

The assignment $G \mapsto \mathrm{B}\,G$ can be made a functor as follows. Let $\phi : G \to H$ be a Lie group homomorphism and let P be a principal H-bundle over X. The fiber bundle $P^\phi = P \times_H G$ associated with P by virtue of the left action of G on H by $(g, h) \mapsto \phi(g)h$ is a principal H-bundle over X, where the action of H is induced from the action of H on itself by right multiplication. Define $\mathrm{B}\,\phi : \mathrm{B}\,G \to \mathrm{B}\,H$ to be the classifying map of the principal H-bundle $(\mathrm{E}\,G)^\phi \to \mathrm{B}\,G$. The assignment $\phi \mapsto \mathrm{B}\,\phi$ is indeed functorial,

$$\mathrm{B}(\psi \circ \phi) = \mathrm{B}\,\psi \circ B\phi\,, \qquad \mathrm{B}\,\mathrm{id}_G = \mathrm{id}_{\mathrm{B}\,G}\,.$$

The classifying map of P^ϕ is $\mathrm{B}\,\phi \circ f : X \to \mathrm{B}\,H$, where $f : X \to \mathrm{B}\,G$ is the classifying map of P. In the special case where ϕ is a Lie subgroup embedding, P^ϕ is usually denoted by P^H and is called the extension of P to the structure group H. Then P is a reduction of P^H to the subgroup G of H. Since the principal G-bundle $\mathrm{E}\,H \to \mathrm{E}\,H/G$ is universal for G, the classifying map $\mathrm{B}\,\phi$ can be identified with the projection in the fiber bundle $\mathrm{E}\,H/G \to \mathrm{B}\,H$ with fiber H/G. For example, the classifying space $\mathrm{B}\,\mathrm{SO}(n)$ is the 2-fold covering of $\mathrm{B}\,\mathrm{O}(n)$.

B Eilenberg-MacLane spaces

Let π be an Abelian group. Up to homotopy equivalence, there exists a unique space X with $\pi_i(X) = \pi$ if $i = n$ and $\pi_i(X) = 0$ otherwise. This space is known as the Eilenberg-MacLane space $K(\pi, n)$. It can be realized as a CW-complex. The following properties of Eilenberg-MacLane spaces are used in the paper:

1. According to the Hurewicz Theorem and the Universal Coefficient Theorem, there is a canonical isomorphism

$$H^n(K(\pi, n), \pi) = \mathrm{Hom}(\pi, \pi)\,.$$

This way, the identity of π is mapped to a distinguished class $\iota_n \in H^n(K(\pi, n), \pi)$, called the fundamental class of $K(\pi, n)$. For any CW-complex X, the assignment of $f^*\iota_n$ to a map $f : X \to K(\pi, n)$ defines a bijection from the set of homotopy classes of maps $X \to K(\pi, n)$ onto the cohomology group $H^n(X, \pi)$.

2. From the long exact homotopy sequence of the path-loop fibration $\Omega K(\pi, n) \hookrightarrow PK(\pi, n) \to K(\pi, n)$ of the space $K(\pi, n)$ one obtains $\Omega K(\pi, n) = K(\pi, n - 1)$.

3. From the long exact homotopy sequence of the product bundle $K(\pi_1, n) \times K(\pi_2, n) \to K(\pi_2, n)$ one obtains $K(\pi_1, n) \times K(\pi_2, n) = K(\pi_1 \oplus \pi_2, n)$.

Furthermore, the notions of transgression and suspension homomorphism are used. Let $F \overset{\iota}{\hookrightarrow} E \xrightarrow{p} B$ be a fibration and $\delta : H^*(F) \to H^*(E, F)$ be the connecting homomorphism of the long exact sequence of the pair (E, F). Let $\tilde{H}^*(B) = H^*(B, *)$ denote the reduced cohomology groups and $\tilde{p} : (E, F) \to (B, *)$ the induced map. Define subsets

$$T^*(F) = \delta^{-1}\tilde{p}^*\tilde{H}^*(B) \subset H^*(F)\,, \qquad S^*(B) = (\tilde{p}^*)^{-1}\delta H^*(F) \subset \tilde{H}^*(B)\,.$$

The transgression is the homomorphism

$$\tau : T^*(F) \to S^*(B)/\ker(\tilde{p}^*)$$

defined by $\tau(u) = [u']$, where $\delta u = \tilde{p}^*u'$. The suspension homomorphism is the homomorphism

$$\sigma : S^*(B) \to T^*(F)/\mathrm{im}\,(\iota^*)$$

defined by $\sigma(v) = [v']$, where $\tilde{p}^*v = \delta v'$. Due to $\ker\tau = \mathrm{im}\,\iota^*$ and $\ker\sigma = \ker\tilde{p}^*$, τ and σ induce inverse isomorphisms

$$T^*(F)/\mathrm{im}\,\iota^* = S^*(B)/\ker\tilde{p}^*.$$

Now consider the fibration $K(\mathbb{Z}_r, n-1) \xrightarrow{\iota} PK(\mathbb{Z}_r, n) \xrightarrow{p} K(\mathbb{Z}_r, n)$. The Serre exact sequence for this fibration contains a portion

$$H^{n-1}(PK(\mathbb{Z}_r, n), \mathbb{Z}_r) \xrightarrow{\iota^*} H^{n-1}(K(\mathbb{Z}_r, n-1), \mathbb{Z}_r) \xrightarrow{\tau} H^n(K(\mathbb{Z}_r, n), \mathbb{Z}_r) \xrightarrow{p^*} H^n(PK(\mathbb{Z}_r, n), \mathbb{Z}_r)\,.$$

Since $H^i(PK(\mathbb{Z}_r, n), \mathbb{Z}_r) = 0$ for all i, $\mathrm{im}\,(\iota^*) = 0 = \mathrm{im}\, p^*$. It follows that τ itself is an isomorphism here and σ is its inverse. Let κ_{n-1} and κ_n denote the fundamental class of $K(\mathbb{Z}_r, n-1)$ and $K(\mathbb{Z}_r, n)$ respectively. Since κ_{n-1} and κ_n are the unique generators of $H^{n-1}(K(\mathbb{Z}_r, n-1), \mathbb{Z}_r)$ and $H^n(K(\mathbb{Z}_r, n), \mathbb{Z}_r)$ respectively, we get

$$\tau(\kappa_{n-1}) = \kappa_n\,, \qquad \sigma(\kappa_n) = \kappa_{n-1}. \tag{76}$$

The definitions of τ and σ given here follow [44]. There is an equivalent definition of the transgression as a certain differential of the Serre spectral sequence. For this as well as for further details we refer to [30].

C Cohomology operations

A cohomology operation is a map $\Theta = \Theta_X : H^m(X, \pi_1) \to H^n(X, \pi_2)$ defined for all spaces X, with a fixed choice of m and n, π_1 and π_2 satisfying the naturality property that for all maps $f : X \to Y$ the following diagram commutes:

$$\begin{tikzcd}
H^m(Y, \pi_1) \arrow[r, "\Theta_Y"] \arrow[d, "f^*"] & H^n(Y, \pi_2) \arrow[d, "f^*"] \\
H^m(X, \pi_1) \arrow[r, "\Theta_X"] & H^n(X, \pi_2)
\end{tikzcd}$$

Let $\iota_m \in H^m(K(\pi_1, m), \pi_1)$ be the fundamental class, see Appendix B. For fixed m, n, π_1 and π_2, the assignment of $\Theta(\iota_m)$ to Θ defines a bijection between the set of all cohomology operations $\Theta : H^m(X, \pi_1) \to H^n(X, \pi_2)$ and the cohomology group $H^n(K(\pi_1, m), \pi_2)$. See e.g. [19, Prop.4I.1] for a proof. The following cohomology operations are used in the text:

1. Let R be a ring. The transformation $H^m(X, R) \to H^{mp}(X, R)$, $\alpha \mapsto \alpha^p$ is a cohomology operation since $f^*(\alpha^p) = (f^*(\alpha))^p$. This example shows that cohomology operations need not be homomorphisms.

2. The Bockstein homomorphisms are cohomology operations. We use in particular the Bockstein homorphisms $\beta_m : H^n(X, \mathbb{Z}_m) \to H^{n+1}(X, \mathbb{Z}_m)$ associated with the coefficient sequence $0 \to \mathbb{Z}_m \overset{m}{\to} \mathbb{Z}_{m^2} \to \mathbb{Z}_m \to 0$ and $\beta : H^n(X, \mathbb{Z}_m) \to H^{n+1}(X, \mathbb{Z})$ associated with the coefficient sequence $0 \to \mathbb{Z} \overset{m}{\to} \mathbb{Z} \overset{\rho_m}{\to} \mathbb{Z}_m \to 0$. They are related via

$$\beta_m = \rho_m \circ \beta\,. \tag{77}$$

We use this relation for the case $m = 2$.

3. The Steenrod square $\mathrm{Sq}^i : H^n(X, \mathbb{Z}_2) \to H^{n+i}(X, \mathbb{Z}_2)$, see e.g. [19], is a cohomology operation uniquely defined by the following axioms:

$$\mathrm{Sq}^0 = \mathrm{id}\,, \tag{78}$$

$$\mathrm{Sq}^i\, x = x^2\,, \quad x \in H^i(X, \mathbb{Z}_2)\,, \tag{79}$$

$$\mathrm{Sq}^i\, x = 0\,, \quad x \in H^j(X, \mathbb{Z}_2)\,,\ j < i\,, \tag{80}$$

$$\mathrm{Sq}^i(x + y) = \mathrm{Sq}^i(x) + \mathrm{Sq}^i(y)\,, \tag{81}$$

$$\mathrm{Sq}^i(xy) = \sum_{j=0}^{i} \mathrm{Sq}^j(x)\, \mathrm{Sq}^{i-j}(y)\,. \tag{82}$$

One has $\mathrm{Sq}^1 = \beta_2$, the Bockstein homomorphism associated with the sequence of coefficient groups $0 \to \mathbb{Z}_2 \xrightarrow{2} \mathbb{Z}_4 \to \mathbb{Z}_2 \to 0$. Hence (77) implies

$$\mathrm{Sq}^1 = \beta_2 = \rho_2 \circ \beta \,. \tag{83}$$

4. The Pontryagin square $\mathfrak{P} : H^{2k}(X, \mathbb{Z}_2) \to H^{4k}(X, \mathbb{Z}_4)$, see [43], is a cohomology operation uniquely defined by the following axioms:

$$\mathfrak{P}\rho u = u^2 \,, \tag{84}$$

$$\rho\mathfrak{P}v = v^2 \,, \tag{85}$$

$$\mathfrak{P}(v_1 + v_2) = \mathfrak{P}v_1 + \mathfrak{P}v_2 + \theta_*(v_1 v_2) \,, \tag{86}$$

for all $u \in H^{4k}(X, \mathbb{Z}_4)$, $v, v_1, v_2 \in H^{2k}(X, \mathbb{Z}_2)$. Writing $u = \rho_4 x$ with $x \in H^{2k}(X, \mathbb{Z})$ in (84) we obtain

$$\mathfrak{P}\rho_2 x = \rho_4 x^2 \,. \tag{87}$$

D Characteristic classes

We collect the basic facts about characteristic classes for principal bundles with structure group $G = \mathrm{O}(n)$, $\mathrm{SO}(n)$ and $\mathrm{Sp}(n)$. For our purposes, it is suitable to view characteristic classes of a G-bundle P as being obtained from generators α of the cohomology of the classifying space $\mathrm{B}\,G$ as

$$\alpha(P) = f^*\alpha \,,$$

where $f : X \to \mathrm{B}\,G$ is the classifying map of P. This way, any relation between generators translates into a relation for characteristic classes of bundles.

Theorem D.1. *$H^*(\mathrm{B\,O}(n), \mathbb{Z}_2)$ is the polynomial ring $\mathbb{Z}_2[w_1, \ldots, w_n]$, with $w_i \in H^i(\mathrm{B\,O}(n), \mathbb{Z}_2)$. Similarly, in the complex case $H^*(\mathrm{B\,U}(n), \mathbb{Z})$ is the polynomial ring $\mathbb{Z}[c_1, \ldots, c_n]$, with $c_i \in H^{2i}(\mathrm{B\,U}(n), \mathbb{Z})$.*

See [20, Thm. 3.2] for a proof. The generators $w_i \in H^i(\mathrm{B\,O}(n), \mathbb{Z}_2)$ are called Stiefel-Whitney classes, the generators $c_i \in H^{2i}(\mathrm{B\,U}(n), \mathbb{Z})$ are called Chern classes.

Theorem D.2.

1. *All torsion elements in $H^*(\mathrm{B\,O}(n), \mathbb{Z})$ are of order 2, and $H^*(\mathrm{B\,O}(n), \mathbb{Z})$ modulo torsion is the polynomial ring $\mathbb{Z}[p_1, \ldots, p_k]$ for $n = 2k$ or $2k+1$.*
2. *All torsion elements in $H^*(\mathrm{B\,SO}(n), \mathbb{Z})$ are of order 2, and $H^*(\mathrm{B\,SO}(n), \mathbb{Z})$ modulo torsion is the polynomial ring $\mathbb{Z}[p_1, \ldots, p_k]$ for $n = 2k+1$ and $\mathbb{Z}[p_1, \ldots, p_{k-1}, e]$ for $n = 2k$, with $e^2 = p_k$ in the latter case.*

Proofs of this theorem can be found in [20] and [19]. The generators p_i are called Pontryagin classes. The generator e is the Euler class of the universal real n-dimensional vector bundle. This class is also present in case $n = 2k+1$ where it is a nontrivial 2-torsion element, uniquely determined by the equation

$$\rho_2 e = w_n \,, \tag{88}$$

see [20, Prop. 3.13]. Application of the long exact sequence associated with the coefficient sequence $0 \to \mathbb{Z} \xrightarrow{2} \mathbb{Z} \xrightarrow{\rho_2} \mathbb{Z}_2 \to 0$ shows that the torsion subgroup of $H^*(\mathrm{B\,O}(n), \mathbb{Z})$ is mapped

injectively to $H^*(\mathrm{B\,O}(n), \mathbb{Z}_2)$, i.e. $H^*(\mathrm{B\,O}(n), \mathbb{Z})$ is completely determined by the classes p_i and w_i. The same argument shows that $H^*(\mathrm{B\,SO}(n), \mathbb{Z})$ is determined by p_i, w_i and e. The injectivity of ρ_2 on torsion elements also follows directly from a Lemma of Borel, see [6], Lemma 24.2.
There exist many relations between the above generators and, hence, between the characteristic classes they define. First, there are relations among the real characteristic classe. As before, let $\beta : H^n(X, \mathbb{Z}_2) \to H^{n+1}(X, \mathbb{Z})$ denote the Bockstein homomorphism associated with the coefficient sequence $0 \to \mathbb{Z} \to \mathbb{Z} \to \mathbb{Z}_2 \to 0$ and let $\rho_m : H^n(X, \mathbb{Z}) \to H^n(X, \mathbb{Z}_m)$ denote the homomorphism induced by reduction mod m.

Theorem D.3. *The following relations hold:*

$$\mathrm{Sq}^j\, w_m = \sum\nolimits_{l=0}^{j} \binom{m-j+l-1}{l} w_{j-l}\, w_{m+l}\,, \tag{89}$$

$$\mathfrak{P} w_{2i+1} = \rho_4 \circ \beta \circ \mathrm{Sq}^{2i}\, w_{2i+1} + \theta_* \left(w_1\, \mathrm{Sq}^{2i}\, w_{2i+1}\right)\,, \tag{90}$$

$$\mathfrak{P} w_{2i} = \rho_4 p_i + \theta_* \left(w_1\, \mathrm{Sq}^{2i-1}\, w_{2i} + \sum\nolimits_{j=0}^{i-1} w_{2j} w_{4i-2j}\right)\,, \tag{91}$$

See [43] and [49] for a proof. Equation (89) is known as the Wu formula.

Proposition D.4. *There holds* $\rho_2 p_i = w_{2i}^2$.

Proof: Consider the coefficient sequence $0 \to \mathbb{Z}_2 \xrightarrow{\theta} \mathbb{Z}_4 \xrightarrow{\rho} \mathbb{Z}_2 \to 0$. By applying ρ to (91) and using $\rho \circ \rho_4 = \rho_2$ and (85) we obtain $0 = \rho \mathfrak{P} w_{2i} - \rho_2 p_i = w_{2i}^2 - \rho_2 p_i$. □

Second, there are relations between real, complex and quaternionic characteristic classes. For clarity, in the following we label the cohomology generators by the groups they belong to. Denote $\tilde{p} = \sum_k (-1)^k p_k$ and $\tilde{c} = \sum_k (-1)^k c_k$. Let

$$\begin{array}{lll} \varphi_{\mathbb{C},\mathbb{R}} : \mathrm{U}(m) \to \mathrm{O}(2m) & \varphi_{\mathbb{H},\mathbb{C}} : \mathrm{Sp}(n) \to \mathrm{U}(2n) & \varphi_{\mathbb{H},\mathbb{R}} = \varphi_{\mathbb{C},\mathbb{R}} \circ \varphi_{\mathbb{H},\mathbb{C}} : \mathrm{Sp}(n) \to \mathrm{O}(4n) \\ j_{\mathbb{R},\mathbb{C}} : \mathrm{O}(n) \to \mathrm{U}(n) & j_{\mathbb{C},\mathbb{H}} : \mathrm{U}(n) \to \mathrm{Sp}(n) & j_{\mathbb{R},\mathbb{H}} : \mathrm{O}(n) \to \mathrm{Sp}(n) \end{array}$$

be the embeddings defined by field restriction and field extension, respectively. The $\varphi_{\cdot,\cdot}$ can be chosen as in Section 3. Recall that the Chern class for $\mathrm{Sp}(n)$-bundles is defined by the element $c(\mathrm{Sp}(n)) := \left(\mathrm{B}\,\varphi_{\mathbb{H},\mathbb{C}}\right)^* c(\mathrm{U}(2n)) \in H^*(\mathrm{B\,Sp}(n), \mathbb{Z})$.

Proposition D.5. *The following relations hold:*

$$\begin{aligned} \left(\mathrm{B}\,\varphi_{\mathbb{C},\mathbb{R}}\right)^* w(\mathrm{SO}(2n)) &= \rho_2\, c(\mathrm{U}(n))\,, \\ \left(\mathrm{B}\,\varphi_{\mathbb{H},\mathbb{R}}\right)^* w(\mathrm{SO}(4n)) &= \rho_2\, c(\mathrm{Sp}(n))\,, \\ \left(\mathrm{B}\,\varphi_{\mathbb{C},\mathbb{R}}\right)^* \tilde{p}(\mathrm{SO}(2n)) &= c(\mathrm{U}(n))\, \tilde{c}(\mathrm{U}(n))\,, \\ \left(\mathrm{B}\,\varphi_{\mathbb{H},\mathbb{R}}\right)^* \tilde{p}(\mathrm{SO}(4n)) &= c(\mathrm{Sp}(n))\, \tilde{c}(\mathrm{Sp}(n))\,, \\ \left(\mathrm{B}\,\varphi_{\mathbb{C},\mathbb{R}}\right)^* e(\mathrm{SO}(2n)) &= c_n(\mathrm{U}(n))\,, \\ \left(\mathrm{B}\,\varphi_{\mathbb{H},\mathbb{R}}\right)^* e(\mathrm{SO}(4n)) &= c_{2n}(\mathrm{Sp}(n))\,. \end{aligned}$$

$$\begin{aligned} (\mathrm{B}\, j_{\mathbb{R},\mathbb{C}})^* c(\mathrm{U}(n)) &= \tilde{p}(\mathrm{O}(n))\,, \\ (\mathrm{B}\, j_{\mathbb{C},\mathbb{H}})^* c(\mathrm{Sp}(n)) &= c(\mathrm{U}(n))\, \tilde{c}(\mathrm{U}(n))\,, \\ (\mathrm{B}\, j_{\mathbb{R},\mathbb{H}})^* c(\mathrm{Sp}(n)) &= \tilde{p}(\mathrm{O}(n))^2\,, \end{aligned}$$

Proof. The relations for $\varphi_{\mathbb{C},\mathbb{R}}$ are well known, see [23], §17 and §20, Proposition 8.4. Since $\varphi_{\mathbb{H},\mathbb{R}} = \varphi_{\mathbb{C},\mathbb{R}} \circ \varphi_{\mathbb{H},\mathbb{C}}$, the relations for $\varphi_{\mathbb{H},\mathbb{R}}$ follow from those for $\varphi_{\mathbb{C},\mathbb{R}}$ by replacing $c(\mathrm{U}(n))$ by $c(\mathrm{Sp}(n))$. The relation for $j_{\mathbb{R},\mathbb{C}}$ is an equivalent definition of Pontryagin classes. To prove the relation for $j_{\mathbb{C},\mathbb{H}}$, observe that by definition of $c(\mathrm{Sp}(n))$, we have

$$(\mathrm{B}\, j_{\mathbb{C},\mathbb{H}})^* c(\mathrm{Sp}(n)) = (\mathrm{B}\, j_{\mathbb{C},\mathbb{H}})^* \,(\mathrm{B}\,\varphi_{\mathbb{H},\mathbb{C}})^*\, c(\mathrm{U}(2n))\,.$$

We compute

$$\varphi_{\mathbb{H},\mathbb{C}} \circ j_{\mathbb{C},\mathbb{H}}(a) = \begin{bmatrix} \overline{a} & 0 \\ 0 & a \end{bmatrix}, \qquad a \in \mathrm{U}(n)\,.$$

Since $c(\mathrm{U}(n)) \mapsto \tilde{c}(\mathrm{U}(n))$ under the map $\mathrm{B}\,\mathrm{U}(n) \to \mathrm{B}\,\mathrm{U}(n)$ induced by conjugation of matrices, $(\mathrm{B}\, j_{\mathbb{C},\mathbb{H}})^* \, (\mathrm{B}\,\varphi_{\mathbb{H},\mathbb{C}})^* \, c(\mathrm{U}(2n)) = c(\mathrm{U}(n))\; \tilde{c}(\mathrm{U}(n))$. To prove the relation for $j_{\mathbb{R},\mathbb{H}}$, we compute

$$(\mathrm{B}\, j_{\mathbb{R},\mathbb{H}})^* c(\mathrm{Sp}(n)) = (\mathrm{B}\, j_{\mathbb{R},\mathbb{C}})^* \, (\mathrm{B}\, j_{\mathbb{C},\mathbb{H}})^* c(\mathrm{Sp}(n)) = (\mathrm{B}\, j_{\mathbb{R},\mathbb{C}})^* \big\{ c(\mathrm{U}(n))\, \tilde{c}(\mathrm{U}(n)) \big\}\,.$$

Due to $(\mathrm{B}\, j_{\mathbb{R},\mathbb{C}})^* \tilde{c}(\mathrm{U}(n)) = \tilde{p}(\mathrm{O}(n))$, the assertion follows. ☐

E The Moore-Postnikov tower

Below we work in the pointed category, hence all maps are assumed to preserve base points. A map $f : X \to Y$ is called an n-equivalence if $f_* : \pi_i(X) \to \pi_i(Y)$ is an isomorphism for $i < n$ and an epimorphism for $i = n$. If f_* is an isomorphism for all i then f is called a weak homotopy equivalence. If $f : X \to Y$ is an n-equivalence and K is a CW-complex then composition with f defines a bijection from the set of homotopy classes of maps $K \to X$ onto the set of homotopy classes of maps $K \to Y$, see e.g. [7, Cor.11.13].

Proposition E.1. *Let $f : X \to Y$ be an $(n+1)$-equivalence and let $F \hookrightarrow X \to Y$ be the corresponding fibration obtained by homotopy equivalent deformation of X. Assume that Y is simply connected. Then the map*

$$f^* : H^k(X, \mathbb{Z}) \to H^k(Y, \mathbb{Z})\,,$$

is an isomorphism for all $k \leq n$, and a monomorphism for $k = n+1$.

Proof: From the exact homotopy sequence of the fibration $F \hookrightarrow X \to Y$ we conclude that $\pi_k(F) = 0$, for $k \leq n$. Using the Hurewicz theorem we get $H_k(F) = \{0\}$ for all $k \leq n$. Inserting this in the universal coefficient theorem

$$H^k(X, \mathbb{Z}) \cong F\big(H_k(X, \mathbb{Z})\big) \oplus \mathrm{Tor}\big(H_{k-1}(X, \mathbb{Z})\big)\,,$$

we obtain $H^k(F, \mathbb{Z}) = \{0\}$ for all $k \leq n$. Since Y is simply connected we obtain (using again the Hurewicz theorem and the universal coefficient theorem) $H^1(Y, \mathbb{Z}) = 0$. Moreover, since Y is simply connected, we have a trivial action of $\pi_1(Y)$ on $H^*(F, \mathbb{Z})$. These facts imply that we have a Serre sequence up to $(n+2)$:

$$\begin{tikzcd} \cdots \arrow[r] & H^{n-1}(F,\mathbb{Z}) \arrow[r] & H^n(Y,\mathbb{Z}) \arrow[r, "f^*"] & H^n(X,\mathbb{Z}) \arrow[r] & H^n(F,\mathbb{Z}) \\ \arrow[r] & H^{n+1}(Y,\mathbb{Z}) \arrow[r, "f^*"] & H^{n+1}(X,\mathbb{Z}) \arrow[r] & H^{n+1}(F,\mathbb{Z}) \arrow[r] & \cdots \end{tikzcd}$$

Since $H^k(F, \mathbb{Z}) = \{0\}$ for all $k \leq n$, the assertion follows. ☐

The Moore-Postnikov-tower of a fibration $F \hookrightarrow E \xrightarrow{p} B$ is a commutative diagram

$$\begin{tikzcd}
 & {} \arrow[d] \\
 & E_n \arrow[d, "p_n"] \\
 & \vdots \arrow[d] \\
 & E_1 \arrow[d, "p_1"] \\
E \arrow[uuur, "q_n"] \arrow[ur, "q_1"] \arrow[r, "p"] & B
\end{tikzcd}$$

such that
1. the maps $q_n : E \to E_n$ are n-equivalences;
2. the maps $P_n = (p_1 \circ p_2 \circ \cdots \circ p_n) : E_n \to B$ induce isomorphisms on π_i for $i > n$ and an injection for $i = n$;
3. the map $p_{n+1} : E_{n+1} \to E_n$ is a fibration with fiber $K(\pi_n(F), n)$.

In Theorem 4.9 we use the convenient fact that every fibration $p : E \to B$ between connected CW-complexes has a Moore-Postnikov tower, see e.g. [19]. For a detailed exposition of Moore-Postnikov theory see [44].

References

[1] Alvarez-Gaumé, L.; Ginsparg, P.: The topological meaning of nonabelian anomalies. Nucl. Phys. **B 243** (1984) 449–474

[2] Asorey, M.; Falceto, F.; López, J.L.; Luzón, G.: Nodes, monopoles, and confinement in $2+1$-dimensional gauge theories. Phys. Lett. B **345** (1995) 125–130

[3] Asorey, M.: Maximal non-Abelian gauges and topology of the gauge orbit space. Nucl. Phys. B **551** (1999) 399–424

[4] Atiyah, M.F.: *K-Theory*. Advanced Book Classics, Addison-Wesley 1989, Notes by D.W. Anderson

[5] Atiyah, M.F.; Singer, I.M.: Dirac operators coupled to vector potentials. Proc. Nat. Acad. Sci. USA **81** (No. 8) (1984) 2597-2600

[6] Borel, A.: *Topics in the Homology Theory of Fibre Bundles*. Lecture Notes in Mathematics 1954, Springer 1967

[7] Bredon, G.E.: *Topology and Geometry*. Springer 1991

[8] Čadek, M.; Vanžura, J.: On oriented vector bundles over CW-complexes of dimension 6 and 7. Comment. Math. Univ. Carolin. **33** , no. 4 (1992) 727–736

[9] Čadek, M.; Vanžura, J.: On the classification of oriented vector bundles over 5-complexes. Czech Math. J. **43** (118) (1993) 753–764

[10] Čadek, M.; Vanžura, J.: On the classification of oriented vector bundles over 9-complexes. Rend. Circ. Mat. Palermo (2) Suppl. (1994), no. 37, 33–40

[11] Carey, A.; Mickelsson, J.; Murray, M.: Index theory, gerbes, and Hamiltonian quantization. Commun. Math. Phys. **183** (1997) 707–722

[12] Charzynski, Sz.; Kijowski, J.; Rudolph, G.; Schmidt, M.: On the stratified classical configuration space of lattice QCD. J. Geom. Phys. **55** (2005) 137–178

[13] Charzynski, Sz.; Rudolph, G.; and Schmidt, M.: On the topology of the reduced classical configuration space of lattice QCD. J. Geom. Phys. **58** (2008) 1607-1623

[14] Emmrich, C.; Römer, H.: Orbifolds as configuration spaces of systems with gauge symmetries. Commun. Math. Phys. **129** (1990) 69–94

[15] Fischer, E.; Rudolph, G.; Schmidt, M.: A lattice gauge model of Singular Marsden-Weinstein Reduction. Part I. Kinematics. J. Geom. Phys. **57** (2007) 1193–1213

[16] Ford, C.; Tok, T.; Wipf, A.: Abelian projection on the torus for general gauge groups. Nucl. Phys. B **548** (1999) 585–612

[17] Ford, C.; Tok, T.; Wipf, A.: SU(N)-gauge theories in Polyakov gauge on the torus. Phys. Lett. B **456** (1999) 155–161

[18] Gribov, V.N.: Quantization of non-Abelian gauge theories. Nucl. Phys. B **139** (1978) 1–19

[19] Hatcher, A.: *Algebraic Topology.* Cambridge University Press, Cambridge, 2002

[20] Hatcher, A.: *Vector Bundles and K-Theory.* http://www.math.cornell.edu∼hatcher/VBKT/VBpage.html

[21] Heil, A.; Kersch, A.; Papadopoulos, N.A.; Reifenhäuser, B.; Scheck, F.: Anomalies from nonfree action of the gauge group. Ann. Phys. **200** (1990) 206–215

[22] Huebschmann, J.; Rudolph, G.; Schmidt, M.: A lattice gauge model for quantum mechanics on a stratified space. hep-th/0702017, to appear in Commun. Math. Phys.

[23] Husemoller, D.: *Fibre Bundles.* Springer 1991

[24] Ireland, K.; Rosen, M.: *A Classical Introduction to Modern Number Theory.* Springer 1990

[25] Kobayashi, S.; Nomizu, K.: *Foundations of Differential Geometry I.* Wiley Classics Library, J. Wiley and Sons 1996

[26] Kondracki, W.; Rogulski, J.: On the notion of stratification. Institute of Mathematics, Polish Academy of Sciences, Preprint 281, Warszawa, 1983

[27] Kondracki, W.; Rogulski, J.: On the stratification of the orbit space for the action of automorphisms on connections. Dissertationes Mathematicae **250**, Panstwowe Wydawnictwo Naukowe, Warszawa, 1986

[28] Langmann, E.; Salmhofer, M.; Kovner, A.: Consistent axial-like gauge fixing on hypertori. Mod. Phys. Lett. A, Vol. 9, No. 31 (1994) 2913–2926

[29] Massey, W.S.: *A Basic Course in Algebraic Topology.* Graduate Texts in Mathematics 127, Springer 1991

[30] McCleary, J.: *A User's Guide to Spectral Sequences.* Cambridge Studies in Advanced Mathematics 58, Cambridge University Press 2001

[31] Milnor, J.: Construction of universal bundles I. Ann. of Math. (2) **63** (1956) 272–284

[32] Milnor, J.: A procedure for killing homotopy groups of differentiable manifolds. Proc. Sympos. Pure Math., Vol. III, American Mathematical Society, Providence, R.I, 1961, pp. 39–55

[33] Narasimhan, M.S.; Ramadas, T.R.: Geometry of SU(2) gauge fields. Commun. Math. Phys. **67** (1979) 121–136

[34] Rudolph, G.; Schmidt, M.; Volobuev, I.P.: Classification of gauge orbit types for SU(n)-gauge theories. Math. Phys. Anal. Geom. **5** (2002) 201–241

[35] Rudolph, G.; Schmidt, M.; Volobuev, I.P.: Partial ordering of gauge orbit types for SU(n)-gauge theories. J. Geom. Phys. **42** (2002) 106–138

[36] Rudolph, G.; Schmidt, M.; Volobuev, I.P.: On the gauge orbit space stratification: a review. J. Phys. A: Math. Gen. **35** (2002) R1–R50

[37] Rudolph, G.; Schmidt, M.: On a certain stratification of the gauge orbit space. Rep. Math. Phys. **50** (2002) 99–110

[38] Schmidt, M.: Classification and partial ordering of reductive Howe dual pairs of classical Lie group. Journal of Geometry and Physics **29** (1999) 283–318

[39] Shabanov, S.V.: 2D Yang Mills theories, gauge orbit spaces and the path integral quantization. Communications in Theoretical Physics (Allahabad), Vol. 4, No. 1 (1995) 1–62

[40] Singer, I.M.: Some remarks on the Gribov ambiguity. Commun. Math. Phys. **60** (1978) 7–12

[41] Steenrod, N.: *The Topology of Fibre Bundles.* Princeton University Press 1951

[42] Thom, R.: Sur une partition en cellules associée à une fonction sur une variété. C. R. Acad. Sci. Paris **228** (1949) 973–975

[43] Thomas, E.: On the cohomology of the real Grassmann complexes and the characteristic classes of n-plane bundles. Trans. Amer. Math. Soc. **96** (1960) 67–89

[44] Thomas, E.: *Seminar on Fiber Spaces.* Lectures delivered in 1964 in Berkeley and 1965 in Zürich. Berkeley notes by J. F. McClendon. Lecture Notes in Mathematics 13, Springer 1966

[45] Thomas, E.: Vector fields on low dimensional manifolds. Math. Z. **103** (1968) 85–93

[46] t'Hooft, G.: On the phase transition towards permanent quark confinement. Nucl. Phys. B **138** (1978) 1–25

[47] Witten, E.: An $SU(2)$-anomaly. Phys. Lett. B **117** (1982) 324–328

[48] Woodward, L.M.: The classification of orientable vector bundles over CW-complexes of small dimension. Proc. Roy. Soc. Edinburgh Sect. A **92** (1982) 175–179

[49] Wu, W.T.: On Pontryagin classes I, II, III. Scientia Sinica **3** (1954) 353–367, Scientia Sinica **4** (1955) 455–490, Acta Math. Sinica **4** (1954) 323–346